



PCT

1
 2
 3
 4
 5
 6
 7
 8
 9
 10
 11
 12
 13
 14
 15
 16
 17
 18
 19
 20
 21
 22
 23
 24
 25
 26
 27
 28
 29
 30
 31
 32
 33
 34
 35
 36
 37
 38
 39
 40
 41
 42
 43
 44
 45
 46
 47
 48
 49
 50
 51
 52
 53
 54
 55
 56
 57
 58
 59
 60
 61
 62
 63
 64
 65
 66
 67
 68
 69
 70
 71
 72
 73
 74
 75
 76
 77
 78
 79
 80
 81
 82
 83
 84
 85
 86
 87
 88
 89
 90
 91
 92
 93
 94
 95
 96
 97
 98
 99
 100
 101
 102
 103
 104
 105
 106
 107
 108
 109
 110
 111
 112
 113
 114
 115
 116
 117
 118
 119
 120
 121
 122
 123
 124
 125
 126
 127
 128
 129
 130
 131
 132
 133
 134
 135
 136
 137
 138
 139
 140
 141
 142
 143
 144
 145
 146
 147
 148
 149
 150
 151
 152
 153
 154
 155
 156
 157
 158
 159
 160
 161
 162
 163
 164
 165
 166
 167
 168
 169
 170
 171
 172
 173
 174
 175
 176
 177
 178
 179
 180
 181
 182
 183
 184
 185
 186
 187
 188
 189
 190
 191
 192
 193
 194
 195
 196
 197
 198
 199
 200
 201
 202
 203
 204
 205
 206
 207
 208
 209
 210
 211
 212
 213
 214
 215
 216
 217
 218
 219
 220
 221
 222
 223
 224
 225
 226
 227
 228
 229
 230
 231
 232
 233
 234
 235
 236
 237
 238
 239
 240
 241
 242
 243
 244
 245
 246
 247
 248
 249
 250
 251
 252
 253
 254
 255
 256
 257
 258
 259
 260
 261
 262
 263
 264
 265
 266
 267
 268
 269
 270
 271
 272
 273
 274
 275
 276
 277
 278
 279
 280
 281
 282
 283
 284
 285
 286
 287
 288
 289
 290
 291
 292
 293
 294
 295
 296
 297
 298
 299
 300
 301
 302
 303
 304
 305
 306
 307
 308
 309
 310
 311
 312
 313
 314
 315
 316
 317
 318
 319
 320
 321
 322
 323
 324
 325
 326
 327
 328
 329
 330
 331
 332
 333
 334
 335
 336
 337
 338
 339
 340
 341
 342
 343
 344
 345
 346
 347
 348
 349
 350
 351
 352
 353
 354
 355
 356
 357
 358
 359
 360
 361
 362
 363
 364
 365
 366
 367
 368
 369
 370
 371
 372
 373
 374
 375
 376
 377
 378
 379
 380
 381
 382
 383
 384
 385
 386
 387
 388
 389
 390
 391
 392
 393
 394
 395
 396
 397
 398
 399
 400
 401
 402
 403
 404
 405
 406
 407
 408
 409
 410
 411
 412
 413
 414
 415
 416
 417
 418
 419
 420
 421
 422
 423
 424
 425
 426
 427
 428
 429
 430
 431
 432
 433
 434
 435
 436
 437
 438
 439
 440
 441
 442
 443
 444
 445
 446
 447
 448
 449
 450
 451
 452
 453
 454
 455
 456
 457
 458
 459
 460
 461
 462
 463
 464
 465
 466
 467
 468
 469
 470
 471
 472
 473
 474
 475
 476
 477
 478
 479
 480
 481
 482
 483
 484
 485
 486
 487
 488
 489
 490
 491
 492
 493
 494
 495
 496
 497
 498
 499
 500
 501
 502
 503
 504
 505
 506
 507
 508
 509
 510
 511
 512
 513
 514
 515
 516
 517
 518
 519
 520
 521
 522
 523
 524
 525

WO 2004/005285 A1

$$\begin{array}{ccc}
 \text{R}_1 & \text{(I)} & \\
 | & & \\
 \text{N}(\text{CH}_2)_p & \text{---} \text{N} & \text{---} \text{C}(=\text{O}) \text{---} \text{R}_2 \\
 & & | \\
 & & \text{O}
 \end{array}
 \qquad
 \begin{array}{ccc}
 \text{R}_6 \text{---} (\text{CH}_2)_n \text{---} \text{A} \text{---} (\text{CH}_2)_m \text{---} \text{N}^+ & \text{(II)} & \\
 | & & \\
 \text{X} & & \text{N}(\text{CH}_2)_p & \text{---} \text{N} & \text{---} \text{C}(=\text{O}) \text{---} \text{R}_2 \\
 & & | & & | \\
 & & \text{O} & & \text{O}
 \end{array}$$

(57) Abstract: New quinuclidine amide derivatives having the chemical structure of general formula (I) and pharmaceutically acceptable salts thereof including quaternary salts of formula (II) are disclosed; as well as processes for their preparation, pharmaceutical compositions 10 comprising them and their use in therapy as antagonists of M3 muscarinic receptors.

QUINUCLIDINE AMIDE DERIVATIVES

This invention relates to new therapeutically useful quinuclidine amide derivatives, to some processes for their preparation and to pharmaceutical compositions containing
5 them.


The novel structures according to the invention are antimuscarinic agents with a potent and long lasting effect. In particular, these compounds show high affinity for M3 muscarinic receptors. This subtype of muscarinic receptor is present in glands and
10 smooth muscle and mediates the excitatory effects of the parasympathetic system on glandular secretion and on the contraction of visceral smooth muscle (Chapter 6, Cholinergic Transmission, in H.P. Rang et al., Pharmacology, Churchill Livingstone, New York, 1995).

15 M3 antagonists are therefore known to be useful for treating diseases characterised by an increased parasympathetic tone, by excessive glandular secretion or by smooth muscle contraction (R.M. Eglen and S.S. Hegde, (1997), Drug News Perspect., 10(8):462-469).

Examples of this kind of diseases are respiratory disorders such as chronic obstructive
20 pulmonary disease (COPD), bronchitis, bronchial hyperreactivity, asthma, cough and rhinitis; urological disorders such as urinary incontinence, pollakiuria, neurogenic or unstable bladder, cystospasm and chronic cystitis; gastrointestinal disorders such as irritable bowel syndrome, spastic colitis, diverticulitis and peptic ulceration; and cardiovascular disorders such as vagally induced sinus bradycardia (Chapter 7,
25 Muscarinic Receptor Agonists and Antagonists, in Goodman and Gilman's The Pharmacological Basis of Therapeutics, 10th edition, McGraw Hill, New York, 2001).

The compounds of the invention can be used alone or in association with other drugs commonly regarded as effective in the treatment of these diseases. For example, they can
30 be administered in combination with β_2 -agonists, steroids, antiallergic drugs, phosphodiesterase IV inhibitors and/or leukotriene D4 (LTD4) antagonists for simultaneous, separate or sequential use in the treatment of a respiratory disease.

The new quinuclidine amide derivatives of the invention have the chemical structure of
35 formula (I):




5 wherein


R_1 represents a hydrogen atom or a straight or branched, optionally substituted lower alkyl group;

10 R_2 represents a group of formula i) or ii)

i)



ii)



wherein

15 R_3 represents a group selected from phenyl, 2-furyl, 3-furyl, 2-thienyl or 3-thienyl;

R_4 represents a group selected from optionally substituted lower alkyl, optionally substituted lower alkenyl, optionally substituted lower alkynyl, cycloalkyl, cycloalkylmethyl, phenyl, benzyl, phenethyl, 2-furyl, 3-furyl, 2-thienyl or 3-thienyl;

20

and R_5 represents a hydrogen atom or a hydroxy, methyl, or $-CH_2OH$ group;

the benzene rings in formula ii) and the cyclic groups represented by R_3 and R_4 being each independently optionally substituted by one, two or three substituents selected from

25 halogen, straight or branched, optionally substituted lower alkyl, hydroxy, straight or branched, optionally substituted lower alkoxy, nitro, cyano, $-CO_2R'$ or $-NR'R''$, wherein R'

and R" each independently represents a hydrogen atom or a straight or branched, optionally substituted lower alkyl group or R' and R" together with the atom to which they are attached form a cyclic group;

5 Q represents a single bond or a -CH₂-, -CH₂-CH₂-, -O-, -O-CH₂-, -S-, -S-CH₂- or -CH=CH- group;

p is 1 or 2 and the amide group is at positions 2, 3 or 4 of the azabicyclic ring;

10 or pharmaceutically acceptable salts thereof; including quaternary ammonium salts;

and all individual stereoisomers and mixtures thereof;


with the proviso that when p is 2, the amide moiety is in position 3 of the quinuclidine ring,

15 R_1 is hydrogen and R_3 and R_4 are both unsubstituted phenyl, then

-when said compound is not a pharmaceutically acceptable salt or is a HCl salt, then R₅ cannot be one of hydrogen or hydroxy; and

20 - when said compound is a quaternary ammonium salt having a methyl group attached to the nitrogen atom of the quinuclidine ring, then R₅ cannot be hydroxy.

The compounds of the invention include quaternary ammonium salts of formula (II).



(11)

25 wherein R_1 , R_2 and p are as defined above;

m is an integer from 0 to 8;

n is an integer from 0 to 4:

30

A represents a group selected from $-\text{CH}_2-$, $-\text{CH}=\text{CR}'-$, $-\text{CR}'=\text{CH}-$, $-\text{CR}'\text{R}''-$, $-\text{C}(\text{O})-$, $-\text{O}-$, $-\text{S}-$, $-\text{S}(\text{O})-$, $-\text{S}(\text{O})_2-$ and $-\text{NR}'-$, wherein R' and R'' are as defined above;

R_6 represents a hydrogen atom, or a group selected from straight or branched, optionally substituted lower alkyl, hydroxy, straight or branched, optionally substituted lower alkoxy, cyano, nitro, $-\text{CH}=\text{CR}'\text{R}''$, $-\text{C}(\text{O})\text{OR}'$, $-\text{OC}(\text{O})\text{R}'$, $-\text{SC}(\text{O})\text{R}'$, $-\text{C}(\text{O})\text{NR}'\text{R}''$, $-\text{NR}'\text{C}(\text{O})\text{OR}''$, $-\text{NR}'\text{C}(\text{O})\text{NR}''$, cycloalkyl, phenyl, naphthanelyl, 5,6,7,8-tetrahydronaphthanelyl, benzo[1,3]dioxolyl, heteroaryl or heterocyclyl; R' and R'' being as defined above; and wherein the cyclic groups represented by R_6 are optionally substituted by one, two or three substituents selected from halogen, hydroxy, straight or branched, optionally substituted lower alkyl, phenyl, $-\text{OR}'$, $-\text{SR}'$, $-\text{NR}'\text{R}''$, $-\text{NHCOR}'$, $-\text{CONR}'\text{R}''$, $-\text{CN}$, $-\text{NO}_2$ and $-\text{COOR}'$; R' and R'' being as defined above;

X^- represents a pharmaceutically acceptable anion of a mono or polyvalent acid.

and all individual stereoisomers and mixtures thereof;

with the proviso that when p is 2, the amide moiety is in position 3 of the quinuclidine ring, R_1 is hydrogen, R_3 and R_4 are both unsubstituted phenyl and R_5 is hydroxy, then in the compounds of formula (II) the sequence $\text{R}_6 - (\text{CH}_2)_n - \text{A} - (\text{CH}_2)_m -$ cannot be a methyl group.

Further objectives of the present invention are to provide processes for preparing said compounds; pharmaceutical compositions comprising an effective amount of said compounds; the use of the compounds in the manufacture of a medicament for the treatment of diseases susceptible of being improved by antagonism of M3 muscarinic receptors; and methods of treatment of diseases susceptible to amelioration by antagonism of M3 muscarinic receptors, which methods comprise the administration of the compounds of the invention to a subject in need of treatment.

Certain quinuclidine amide derivatives, which fall outside the scope of the present invention, have been disclosed in JP 09328469 and WO 9316048.

As used herein, an alkyl group or moiety can be straight or branched, and is typically a lower alkyl group or moiety. A lower alkyl group or moiety contains 1 to 8, preferably 1 to 6

and more preferably 1 to 4 carbon atoms. In particular it is preferred that such an alkyl group or moiety is represented by a methyl, ethyl, propyl, including i-propyl or butyl, including n-butyl, sec-butyl and tert-butyl group or moiety.

- 5 As used herein, an alkenyl group or moiety can be straight or branched, and is typically a lower alkenyl group or moiety. A lower alkenyl group or moiety contains 2 to 8, preferably 2 to 6 and more preferably 2 to 4 carbon atoms. In particular it is preferred that such an alkenyl group or moiety is represented by a vinyl, allyl or 1-propenyl group or moiety.
- 10 As used herein, an alkynyl group or moiety can be straight or branched, and is typically a lower alkynyl group or moiety. A lower alkynyl group or moiety contains 2 to 8, preferably 2 to 6 and more preferably 2 to 4 carbon atoms. In particular it is preferred that such an alkynyl group or moiety is represented by a 3-butyne or 1-propyne group or moiety.
- 15 Optionally substituted lower alkyl, alkenyl or alkynyl groups mentioned herein include straight or branched lower alkyl, alkenyl or alkynyl groups as defined above, which may be unsubstituted or substituted in any position by one or more substituents, for example by 1, 2 or 3 substituents. When two or more substituents are present, each substituent may be the same or different. The substituent(s) are typically halogen atoms, preferably fluoride
- 20 atoms, and hydroxy or alkoxy groups.

- Alkoxy groups mentioned herein are typically lower alkoxy groups, that is groups containing from 1 to 8, preferably 1 to 6 and more preferably 1 to 4 carbon atoms, the hydrocarbon chain being branched or straight and optionally substituted in any position by
- 25 one or more substituents, for example by 1, 2 or 3 substituents. When two or more substituents are present, each substituent may be the same or different. The substituent(s) are typically halogen atoms, most preferably fluoride atoms, and hydroxy groups. Preferred optionally substituted alkoxy groups include methoxy, ethoxy, n-propoxy, i-propoxy, n-butoxy, sec-butoxy, t-butoxy, trifluoromethoxy and difluoromethoxy,
- 30 hydroxymethoxy, hydroxyethoxy or 2-hydroxypropoxy.

Cyclic groups mentioned herein include carbocyclic and heterocyclic groups. The cyclic groups can contain one or more rings. Carbocyclic groups may be aromatic or alicyclic, for example cycloalkyl groups. Heterocyclic groups also include heteroaryl groups.

Cycloalkyl groups and alicyclic groups mentioned herein, unless otherwise specified, typically contain from 3 to 7 carbon atoms. Cycloalkyl groups and alicyclic rings of 3 to 7 carbon atoms include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl.

- 5 As used herein an aromatic group typically contains from 5 to 14, preferably 5 to 10 carbon atoms. Examples of aromatic groups include phenyl and naphthalenyl.

- A heterocyclic or heteroaromatic group mentioned herein is typically a 5 to 10 membered group, such as a 5, 6 or 7 membered group, containing one or more heteroatoms selected from N, S and O. Typically, 1, 2, 3 or 4 heteroatoms are present, preferably 1 or 2 heteroatoms. A heterocyclic or heteroaromatic group may be a single ring or two or more fused rings wherein at least one ring contains a heteroatom. Examples of heterocyclic groups include piperidyl, pyrrolidyl, piperazinyl, morpholinyl, thiomorpholinyl, pyrrolyl, imidazolyl, imidazolidinyl, pyrazolyl, indolyl, isoindolyl, pyridyl, pyrazinyl, pyrimidinyl, 10 pyridazinyl, indolizyl, isoindolyl, indolyl, indazolyl, purinyl, quinolizyl, isoquinolyl, quinolyl, phthalazinyl, naphthyridinyl, quinoxalyl, quinazolyl, cinnolyl, pteridinyl, quinuclidinyl, triazolyl, pyrazolyl, tetrazolyl, thienyl and dioxolyl. Examples of heteroaromatic groups include pyridyl, thienyl, furyl, pyrrolyl, imidazolyl, benzothiazolyl, pyridinyl, pyrazolyl, pyrazinyl, pyrimidinyl, pyridazinyl, indolyl, indazolyl, purinyl, quinolyl, 15 isoquinolyl, phthalazinyl, naphthyridinyl, quinoxalyl, quinazolyl, cinnolyl, triazolyl and pyrazolyl.

- As used herein a halogen atom includes a fluorine, chlorine, bromine or iodine atom, typically a fluorine, chlorine or bromine atom.

25

- As used herein, a pharmaceutically acceptable salt is a salt with a pharmaceutically acceptable acid or base. Pharmaceutically acceptable acids include both inorganic acids, for example hydrochloric, sulphuric, phosphoric, diphosphoric, hydrobromic and nitric acid and organic acids, for example citric, fumaric, formic, maleic, malic, ascorbic, succinic, 30 tartaric, benzoic, acetic, methanesulphonic, ethanesulphonic, benzenesulphonic or p-toluenesulphonic acid. Pharmaceutically acceptable bases include alkali metal (e.g. sodium or potassium) and alkali earth metal (e.g. calcium or magnesium) hydroxides and organic bases, for example alkyl amines, aralkyl amines and heterocyclic amines. Preferred salts of the compounds of formula (I) are those formed between the nitrogen 35 atom of the azabicyclic ring and an inorganic or an organic acid

Other preferred salts according to the invention are quaternary ammonium compounds of formula (II), wherein an equivalent of an anion (X^-) is associated with the positive charge of the N atom. X^- may be an anion of various mineral acids such as, for example, chloride, bromide, iodide, sulphate, nitrate, phosphate, or an anion of an organic acid such as, for example, acetate, formate, maleate, fumarate, citrate, oxalate, succinate, tartrate, malate, mandelate, trifluoroacetate, methanesulphonate and p-toluenesulphonate. X^- is preferably an anion selected from chloride, bromide, iodide, sulphate, nitrate, acetate, maleate, oxalate, succinate or trifluoroacetate. More preferably X^- is chloride, bromide, trifluoroacetate or methanesulphonate.

Preferred compounds of formulae (I) or (II) are those wherein R_1 is hydrogen, methyl or ethyl, most preferably hydrogen.

In some preferred embodiments of the invention R_2 is a group of formula i), wherein R_3 is a group, which is optionally substituted with one or more halogen atom(s), selected from phenyl, 2-thienyl, 3-thienyl or 2-furyl. More preferably R_2 is a group of formula i), wherein R_3 represents a group phenyl, 2-thienyl or 2-furyl which are optionally substituted with one or more halogen atom(s). Most preferably, R_3 is phenyl or 2-thienyl,

20

Further preferred compounds having a group of formula i) are those wherein R_4 represents a linear group selected from ethyl, n-butyl, vinyl, allyl, 1-propenyl and 1-propynyl, or a group, which is optionally substituted with one or more halogen atom(s), methyl or methoxy group(s), selected from cyclopentyl, cyclohexyl, phenyl, benzyl,

phenethyl, 2-thienyl and 3-furyl. More preferably compounds having a group of formula i) are those wherein R_4 represents a linear group selected from ethyl, n-butyl, vinyl, allyl and 1-propynyl, or a group, which is optionally substituted with one or more halogen atom(s), methyl or methoxy group(s), selected from cyclopentyl, phenyl, benzyl, phenethyl and 2-thienyl. Most preferably, R_4 is ethyl, n-butyl, vinyl, allyl, cyclopentyl, phenyl, benzyl or 2-thienyl.

30

In other preferred embodiments of the invention R_2 is a group of formula ii), wherein Q represents a single bond or an oxygen atom.

In the most preferred embodiments of the invention R_5 in groups i) or ii) is a hydrogen atom or a hydroxy group.

Preferably, in the compounds of formulae (I) or (II) p is 2 and the amide group is at
5 positions 3 or 4 of the azabicyclic ring, most preferably at position 3.

Particularly preferred compounds of formula (II) are those wherein m is an integer from 0 to 6 and n is an integer from 0 to 4; most preferably m is 0 to 5 and n 0 to 2; A represents a group selected from $-CH_2-$, $-CH=CH-$, $-O-$, $-C(O)-$, $-NR'-$, and $-S-$; most preferably A is –
10 CH_2- , $-CH=CH-$, $-O-$; and R_6 is a hydrogen atom, a cyano group, a nitro group, a –
 $C(O)OR'$ group, a $-OC(O)R'$ group, a $-SC(O)R'$ group, a $-CH=CH_2$ group, a $-CH=CR'R''$ group, a $C(O)NR'R''$ group, a straight or branched C_1-C_4 alkyl group, which is optionally substituted with one or more halogen atom(s), a straight C_1-C_4 alkoxy group, which is optionally substituted with one or more halogen atom(s) or hydroxy group(s), or a cyclic
15 group, which is optionally substituted with one or more substituents selected from halogen atoms, groups of formula $-C(O)NR'R''$ or methyl, hydroxy, nitro or phenyl groups, the cyclic group being selected from cyclohexyl, phenyl, 5,6,7,8-tetrahydronaphthanelyl, 2-thienyl, 1-pyrrolidiny, 1-pyrrolyl, benzo[1,3]dioxolyl, 2-benzothiazolyl, naphthalenyl and dioxolyl. More preferably compounds of formula (II) are those wherein m is an integer
20 from 0 to 5 and n is an integer from 0 to 2; A represents a group selected from $-CH_2-$, $-CH=CH-$, $-O-$, $-C(O)-$, $-NR'-$, and $-S-$; and R_6 is a hydrogen atom, a cyano group, a –
 $C(O)OR'$ group, a $-OC(O)R'$ group, a $-SC(O)R'$ group, a $-CH=CH_2$ group, a $-C(O)NR'R''$ group, a straight or branched C_1-C_4 alkyl group, a trifluoromethyl, or a cyclic group selected from cyclohexyl, 5,6,7,8-tetrahydronaphthanelyl, 2-thienyl, 1-pyrrolyl,
25 benzo[1,3]dioxolyl, 2-benzothiazolyl, naphthalenyl, dioxolyl and phenyl, which is optionally substituted with one or more substituents selected from halogen atoms, groups of formula $-C(O)NR'R''$, methyl, hydroxy and phenyl groups. Most preferably R_6 is selected from hydrogen, straight C_1-C_4 alkyl, $-CH=CH_2$, cyclohexyl, phenyl which is unsubstituted or substituted with one or two substituents selected from methyl groups and
30 hydroxy groups, 5,6,7,8-tetrahydronaphthanelyl and 2-thienyl.

Most preferred are compounds of formula (II) wherein the sequence $R_6 - (CH_2)_n - A - (CH_2)_m -$ is one of methyl, 3-phenoxypropyl, 3-(3-hydroxyphenoxy)propyl, allyl, heptyl, 3-phenylpropyl, 3-phenylallyl, 2-phenoxyethyl, 2-benzyloxyethyl, cyclohexylmethyl, 3-

(5,6,7,8-tetrahydronaphthalen-2-yloxy)propyl, 5-(2,6-dimethylphenoxy)pentyl, 3-thien-2-ylpropyl or 3-cyclohexylpropyl and X⁻ is bromide or trifluoroacetate.

The compounds of the present invention represented by formulae (I) or (II) described
 5 above may have one or more asymmetric atoms, e.g. the carbon in position 3 of the quinuclidine ring; the carbon substituted by R₃, R₄ and R₅ in the compounds wherein R₂ is a group of formula i); or the carbon which is attached to the carbonyl group in the compounds wherein R₂ is a group of formula ii). Each of these asymmetric atoms may have R- or S-configuration. The single isomers and mixtures of the isomers fall within the
 10 scope of the invention.

Particular individual compounds of formula (I) include:

- N-(1-Azabicyclo[2.2.2]oct-3-yl)-2-hydroxy-2,2-dithien-2-ylacetamide
- 15 N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2,2-dithien-2-ylacetamide
 N-[(3S)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2,2-dithien-2-ylacetamide
 N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2,2-dithien-2-ylacetamide
 N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-cyclopentyl-2-hydroxy-2-thien-2-ylacetamide
 N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2-thien-2-ylpent-4-enamide
- 20 (2*)-N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2-thien-2-ylbutanamide
 (diastereomer 1)
 (2*)-N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2-thien-2-ylbutanamide
 (diastereomer 2)
 (2*)-N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2-thien-2-ylbut-3-enamide
- 25 (diastereomer 1)
 (2*)-N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2-thien-2-ylbut-3-enamide
 (diastereomer 2)
 N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2,3-diphenylpropanamide
 N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-cyclopentyl-2-hydroxy-2-phenylacetamide
- 30 N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-phenylhexanamide
 N-(1-Azabicyclo[2.2.2]oct-3-yl)-9H-xanthene-9-carboxamide
 N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-9H-xanthene-9-carboxamide
 N-[(3S)-1-Azabicyclo[2.2.2]oct-3-yl]-9H-xanthene-9-carboxamide
 N-(1-Azabicyclo[2.2.2]oct-3-yl)-9-hydroxy-9H-fluorene-9-carboxamide
- 35 N-1-Azabicyclo[2.2.2]oct-3-yl-N-methyl-9H-xanthene-9-carboxamide

- N-1-Azabicyclo[2.2.2]oct-3-yl-N-ethyl-9H-xanthene-9-carboxamide
 N-1-Azabicyclo[2.2.2]oct-4-yl-2-cyclopentyl-2-hydroxy-2-thien-2-ylacetamide
 N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2-phenyl-2-thien-2-ylacetamide
 N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-(5-bromothien-2-yl)-2-(4-fluoro-3-methylphenyl)-2-
 5 hydroxyacetamide
 N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-fur-2-yl-2-hydroxypent-3-ynamide
 N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-fur-2-yl-2-hydroxy-4-(4-methoxyphenyl)butanamide
 N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-fur-3-yl-2-hydroxy-2-thien-2-ylacetamide
 N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-cyclohexyl-2-hydroxy-2-thien-3-ylacetamide
 10 N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-9-hydroxy-9H-fluorene-9-carboxamide
 N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-N-methyl-9H-xanthene-9-carboxamide
 (2S)-N-[(3R)-1-azabicyclo[2.2.2]oct-3-yl]-2-cyclopentyl-2-hydroxy-2-thien-2-ylacetamide

- ((*) Configuration not assigned; either the (2R)- or the (2S)- isomers of the above
 15 compounds may be produced).

Particular individual compounds of formula (II) include:

- 3-(2-Hydroxy-2,2-dithien-2-ylacetyl-amino)-1-methyl-1-azoniabicyclo[2.2.2]octane bromide
 20 3-(2-Hydroxy-2,2-dithien-2-ylacetyl-amino)-1-(3-phenoxypropyl)-1-
 azoniabicyclo[2.2.2]octane bromide
 (3R)-3-(2-Hydroxy-2,2-dithien-2-ylacetyl-amino)-1-methyl-1-azoniabicyclo[2.2.2]octane
 trifluoroacetate
 (3R)-1-Allyl-3-(2-hydroxy-2,2-dithien-2-ylacetyl-amino)-1-azoniabicyclo[2.2.2]octane
 25 trifluoroacetate
 (3R)-1-Heptyl-3-(2-hydroxy-2,2-dithien-2-ylacetyl-amino)-1-azoniabicyclo[2.2.2]octane
 trifluoroacetate
 (3R)-3-(2-Hydroxy-2,2-dithien-2-ylacetyl-amino)-1-(3-phenylpropyl)-1-
 azoniabicyclo[2.2.2]octane bromide
 30 (3R)-3-(2-Hydroxy-2,2-dithien-2-ylacetyl-amino)-1-((E)-3-phenylallyl)-1-
 azoniabicyclo[2.2.2]octane trifluoroacetate
 (3R)-3-(2-Hydroxy-2,2-dithien-2-ylacetyl-amino)-1-(2-phenoxyethyl)-1-
 azoniabicyclo[2.2.2]octane bromide
 (3R)-3-(2-Hydroxy-2,2-dithien-2-ylacetyl-amino)-1-(3-phenoxypropyl)-1-
 35 azoniabicyclo[2.2.2]octane bromide

- (3R)-3-(2-Hydroxy-2,2-dithien-2-ylacetyl amino)-1-[3-(3-hydroxyphenoxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-1-(2-Benzoyloxyethyl)-3-(2-hydroxy-2,2-dithien-2-ylacetyl amino)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 5 (3R)-3-(2-Hydroxy-2,2-dithien-2-ylacetyl amino)-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane bromide
- (3S)-3-(2-Hydroxy-2,2-dithien-2-ylacetyl amino)-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane bromide
- (3R)-3-(2,2-Dithien-2-ylacetyl amino)-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane
- 10 bromide
- 1-Methyl-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane bromide
- 1-(3-Phenoxypropyl)-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane bromide
- (3R)-1-(3-Phenoxypropyl)-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane bromide
- 15 (3S)-1-Allyl-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3S)-1-Heptyl-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 20 (3S)-1-Cyclohexylmethyl-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3S)-1-(3-Cyclohexylpropyl)-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3S)-1-(3-Phenoxypropyl)-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane bromide
- 25 (3S)-1-[3-(5,6,7,8-Tetrahydronaphthalen-2-yloxy)propyl]-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3S)-1-[5-(2,6-Dimethylphenoxy)pentyl]-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 30 3-[(9-Hydroxy-9H-fluoren-9-yl)carbonyl]amino-1-methyl-1-azoniabicyclo[2.2.2]octane bromide
- 3-[(9-Hydroxy-9H-fluoren-9-yl)carbonyl]amino-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane bromide
- (3R)-1-[3-(2-Carbamoylphenoxy)propyl]-3-[(9-Hydroxy-9H-fluoren-9-yl)carbonyl]amino-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 35


- (3R)-1-[4-(4-Fluorophenyl)-4-oxobutyl]-3-[(9-Hydroxy-9H-fluoren-9-yl)carbonylamino]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(9-Hydroxy-9H-fluoren-9-yl)carbonylamino]-1-(4-oxo-4-thien-2-ylbutyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 5 (3R)-3-[(9-Hydroxy-9H-fluoren-9-yl)carbonylamino]-1-[3-(methylphenylamino)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[(9-Hydroxy-9H-fluoren-9-yl)carbonylamino]-1-(3-phenylsulfanylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 3-[Methyl-(9H-xanthen-9-ylcarbonyl)amino]-1-(3-pyrrol-1-ylpropyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 10 1-[3-(Biphenyl-4-yloxy)propyl]-3-[ethyl-(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-1-[3-(Benzo[1,3]dioxol-5-yloxy)propyl]-3-(2-hydroxy-2-phenyl-2-thien-2-ylacetyl-amino)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 15 (3R)-1-[3-(Benzothiazol-2-yloxy)propyl]-3-(2-hydroxy-2-phenyl-2-thien-2-ylacetyl-amino)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-(2-Fur-2-yl-2-hydroxypent-3-ynoylamino)-1-[3-(naphthalen-1-yloxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-[2-Fur-2-yl-2-hydroxy-4-(4-methoxyphenyl)butyrylamino]-1-[6-(4-phenylbutoxy)hexyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 20 (3R)-3-(2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetyl-amino)-1-(2-hydroxyethyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-(2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetyl-amino)-1-(2-ethoxyethyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 25 (3R)-3-(2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetyl-amino)-1-{2-[2-(2-hydroxyethoxy)ethoxy]ethyl}-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-(2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetyl-amino)-1-(4,4,4-trifluorobutyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-1-(4-Acetoxybutyl)-3-[2-(5-bromothien-2-yl)-2-(4-fluoro-3-methylphenyl)-2-hydroxyacetyl-amino]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 30 (3R)-3-[2-(5-Bromothien-2-yl)-2-(4-fluoro-3-methylphenyl)-2-hydroxyacetyl-amino]-1-(4-ethoxycarbonylbutyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-1-(3-Acetylsulfanylpropyl)-3-[2-(5-bromothien-2-yl)-2-(4-fluoro-3-methylphenyl)-2-hydroxyacetyl-amino]-1-azoniabicyclo[2.2.2]octane trifluoroacetate

- (3R)-1-(3-Cyanopropyl)-3-(2-hydroxy-2-phenyl-2-thien-2-ylacetyl-amino)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-1-(2-Carbamoylethyl)-3-(2-hydroxy-2-thien-2-ylpent-4-enoylamino)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 5 (3R)-1-(2-[1,3]Dioxolan-2-ylethyl)-3-(2-hydroxy-2-thien-2-ylpent-4-enoylamino)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 4-(2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetyl-amino)-1-(4-methylpent-3-enyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-1-[3-(2-Carbamoylphenoxy)propyl]-3-[[9-Hydroxy-9H-fluoren-9-yl]carbonyl]amino)-1-azoniabicyclo[2.2.2]octane formate
- 10 (3R)-1-[4-(4-Fluorophenyl)-4-oxobutyl]-3-[[9-Hydroxy-9H-fluoren-9-yl]carbonyl]amino)-1-azoniabicyclo[2.2.2]octane formate
- (3R)-3-[[9-Hydroxy-9H-fluoren-9-yl]carbonyl]amino)-1-[3-(methylphenylamino)propyl]-1-azoniabicyclo[2.2.2]octane chloride
- 15 (3R)-3-[[9-Hydroxy-9H-fluoren-9-yl]carbonyl]amino)-1-(3-phenylsulfanylpropyl)-1-azoniabicyclo[2.2.2]octane formate
- (3R)-3-[Methyl-(9H-xanthen-9-ylcarbonyl)amino]-1-(3-pyrrol-1-ylpropyl)-1-azoniabicyclo[2.2.2]octane bromide
- (3R)-1-[3-(Biphenyl-4-yloxy)propyl]-3-[methyl-(9H-xanthene-9-carbonyl)amino]-1-azoniabicyclo[2.2.2]octane chloride
- 20 (3R)-3-(2-Fur-2-yl-2-hydroxypent-3-ynoylamino)-1-[3-(naphthalen-1-yloxy)propyl]-1-azoniabicyclo[2.2.2]octane chloride
- (3R)-1-[3-(Benzo[1,3]dioxol-5-yloxy)propyl]-3-(2-fur-2-yl-2-hydroxypent-3-ynoylamino)-1-azonia-bicyclo[2.2.2]octane bromide
- 25 (3R)-1-[3-(Benzothiazol-2-yloxy)propyl]-3-(2-fur-2-yl-2-hydroxypent-3-ynoylamino)-1-azonia-bicyclo[2.2.2]octane chloride
- (3R)-3-[[2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetyl]amino)-1-(2-hydroxyethyl)-1-azoniabicyclo[2.2.2]octane bromide
- (3R)-3-[[2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetyl]amino)-1-(2-ethoxyethyl)-1-azoniabicyclo[2.2.2]octane formate
- 30 (3R)-3-[[2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetyl]amino)-1-(4,4,4-trifluorobutyl)-1-azoniabicyclo[2.2.2]octane bromide
- (3R)-1-(4-Acetoxybutyl)-3-[2-(5-bromothien-2-yl)-2-(4-fluoro-3-methylphenyl)-2-hydroxyacetyl-amino]-1-azoniabicyclo[2.2.2]octane bromide

- (3R)-3-[2-(5-Bromothien-2-yl)-2-(4-fluoro-3-methylphenyl)-2-hydroxyacetyl-amino]-1-(4-ethoxycarbonylbutyl)-1-azoniabicyclo[2.2.2]octane bromide
- (3R)-1-(3-Acetylsulfanylpropyl)-3-[2-(5-bromothien-2-yl)-2-(4-fluoro-3-methylphenyl)-2-hydroxyacetyl-amino]-1-azoniabicyclo[2.2.2]octane formate
- 5 (3R)-1-(3-Cyanopropyl)-3-[2-fur-2-yl-2-hydroxy-4-(4-methoxyphenyl)butyrylamino]-1-azoniabicyclo[2.2.2]octane bromide
- (3R)-1-(2-Carbamoylethyl)-3-[2-fur-2-yl-2-hydroxy-4-(4-methoxyphenyl)butyrylamino]-1-azoniabicyclo[2.2.2]octane formate
- (3R)-1-(2-[1,3]Dioxolan-2-yl-ethyl)-3-[2-fur-2-yl-2-hydroxy-4-(4-
- 10 methoxyphenyl)butyrylamino]-1-azoniabicyclo[2.2.2]octane bromide

In accordance with another embodiment, the present invention provides processes for preparing the novel quinuclidine amide derivatives of formulae (I) and (II)

- 15 The compounds of formula (I) may be prepared from the corresponding carboxylic acid of formula (III) following two different methods (a) and (b), illustrated in the following scheme.




20

Some compounds of formula (IV) are commercially available, for example 3-aminoquinuclidine (Aldrich, dihydrochloride), (3R)-aminoquinuclidine (Aldrich, Finorga, dihydrochloride) and (3S)-aminoquinuclidine (Aldrich, dihydrochloride).

(3R)-N-methylquinuclidin-3-amine has been prepared as described in the experimental section.

- 5 Those compounds of formula (I), wherein R_2 is a group of formula i) and R_5 is an hydroxy group may also be prepared from the compounds of formula (V) by reaction with the corresponding organometallic derivative.



10

The compounds of formula (V) may be prepared from the corresponding glyoxylic acids following methods equivalent to methods (a) and (b) described in Scheme 1

15 Certain compounds of formula (V) are novel and fall within the scope of the present invention. In particular:

N-1-azabicyclo[2.2.2]oct-3-yl-2-oxo-2-thien-2-ylacetamide


N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-oxo-2-thien-2-ylacetamide

N-[(3S)-1-Azabicyclo[2.2.2]oct-3-yl]-2-oxo-2-thien-2-ylacetamide

20


The quaternary ammonium derivatives of formula (II) may be prepared, as illustrated in the following scheme, by reaction of the compounds of formula (I) with an alkylating agent of formula (VI) using two possible methods -(d) or (e)-, described in detail in the experimental section. Method (e) involves the use of solid phase extraction techniques

25 that allow the parallel preparation of several compounds.



In formula (VI), W represents any suitable leaving group, preferably a group X^- as defined above for the compounds of formula (II). When W is a leaving group other than X^- , the quaternary ammonium salt of formula (II) is produced from the product of method (d) or (e) by an exchange reaction according to standard methods to replace the anion W with the desired anion X^- .

Those compounds of general formula (VI) which are not commercially available have been prepared according to standard methods. For example, compounds wherein $n=0$ and $\text{A} = -\text{O}-, -\text{S}-, \text{NR}'$ were obtained by reaction of the corresponding alcohol, thiol or amine derivative, or its sodium or potassium salt, with an alkylating agent of general formula $\text{Y}-(\text{CH}_2)_m-\text{W}$, wherein W is as defined above; most preferably W is a halogen atom and Y a halogen atom or a sulphonate ester. In other examples, compounds of general formula (VI) wherein n is at least 1 were synthesised from the corresponding alcohol derivative of general formula (VII) by methods known in the art.



The compounds of formulae (I) and (II) can have one or more asymmetric carbons. All possible stereoisomers, single isomers and mixtures of isomers are also included within

the scope of the present invention. The diastereomers of the compounds may be separated by conventional methods, for example by chromatography or crystallisation.

The structures of the compounds were confirmed by ^1H -NMR and MS. The NMR spectra were recorded using either a Varian 300 MHz instrument or a Bruker DPX-250 instrument. Chemical shifts are expressed as parts per million (δ) from the internal reference tetramethylsilane. The purity of the compounds was determined by HPLC, using reverse phase chromatography on a Waters instrument. Molecular ions were produced by electrospray ionisation mass spectrometry on a Hewlett Packard instrument.

10

Method (a)

Preparation of N-[(3R)-1-azabicyclo[2.2.2]oct-3-yl]-2-phenylhexanamide (Example 27)

15 2-phenylhexanoic acid was prepared by alkylation of phenylacetic acid with 1-chlorobutane following a standard method. Oxalyl chloride (0.88 ml, 0.0101 mol) was then added to a solution of 1.62 g of 2-phenylhexanoic acid (0.0084 mol) and dimethylformamide (DMF, one drop) in 25 ml of CHCl_3 (ethanol free) at 0°C . The mixture was stirred and allowed to warm at room temperature. After an hour at this temperature
20 the solvents were evaporated and the residue was dissolved in CHCl_3 and evaporated again. This procedure was repeated two times. The oil obtained was dissolved in CHCl_3 and the solution obtained cooled at 0°C . A solution of (3R)-aminoquinuclidine (1.28 g, 0.0101 mol) in CHCl_3 was added. The mixture was stirred and allowed to warm at room temperature. After 1 hour at this temperature, the reaction mixture was diluted with CHCl_3
25 and washed with a 10% solution of aqueous potassium carbonate; then washed with water, dried over Na_2SO_4 and evaporated to give 3.57 g of an oil which was purified by column chromatography (silica gel, CHCl_3 :MeOH: NH_4OH 90:10:1 as eluent). Appropriate fractions were combined and evaporated to give 2.42 g of a solid that after treatment with isopropyl ether yielded 1.59 g (63.1%) of the title product as a mixture of
30 diastereomers.

mp: 136°C .

MS $[\text{M}+1]^+$: 301

^1H -NMR ($\text{DMSO}-d_6$): (mixture of diastereomers 50:50) δ 0.84 (t, 3H), 1.10-1.60 (m, 9H), 1.70 (m, 1H), 1.90 (m, 1H), 2.25-2.50 (m, 1H), 2.50-2.80 (m, 4H), 2.94 & 3.05 (m,
35 1H), 3.50 (m, 1H), 3.65 (m, 1H), 7.15-7.40 (m, 5H), 8.04 (m, 1H, NH).

Preparation of N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-N-methyl-9H-xanthene-9-carboxamide (Example 47)

Oxalyl chloride (0.540 ml, 0.0062 mol) was added to a solution of 9H-xanthene-9-carboxylic acid (1.16 g, 0.0051 mol) and dimethylformamide (two drops) in 20 ml of CHCl_3 (ethanol free) at 0°C . The mixture was allowed to warm to room temperature under stirring and maintained 1 h at this temperature. After this time, the reaction mixture was concentrated to dryness in vacuo and the obtained residue was dissolved in CHCl_3 (15 ml) and concentrated again. This procedure was repeated two times. The obtained residue was dissolved in CHCl_3 and the solution cooled to 0°C . A solution of 0.865 g (0.0062 mol) of (3R)-N-methylquinuclidin-3-amine (Intermediate I-4) in 10 ml of CHCl_3 was added. The mixture was allowed to warm to room temperature under stirring. After 4 hours at this temperature, the reaction mixture was treated with K_2CO_3 (saturated solution) and the aqueous phase was extracted with CHCl_3 . The organic solutions were combined and washed with K_2CO_3 (saturated solution) and water, dried over Na_2SO_4 filtered and concentrated to dryness. The obtained residue was purified by column chromatography (silica gel, CHCl_3 : NH_4OH 99:1 \rightarrow CHCl_3 : MeOH : NH_4OH 98:2:1 as eluent) to yield 900 mg (50%) of the title product.

MS $[\text{M}+1]^+$: 349

^1H -RMN (400 MHz, 60°C , CDCl_3): δ 7.26 - 7.19 (m, 4H), 7.10 - 7.01 (m, 4H), 5.51 (s, 1H), 4.18 (broad multiplet, 1H), 2.92 (s, 3H), 2.85-2.40 (m, 6H), 1.85-1.20 (m, 5H).

Preparation of (3R)-N-methylquinuclidin-3-amine (Intermediate I-4)

1g (0.0079 mol) of (3R)-aminoquinuclidine was dissolved in 20 ml of CH_2Cl_2 and 1.22 ml (0.0087 mol) of triethylamine were added. The obtained solution was cooled to 0°C , ClCO_2Et (0.835 ml, 0.0087 mol) was added and the mixture was stirred 15 h at room temperature. After this time, solvents were evaporated in vacuo, the obtained residue was dissolved in CHCl_3 and the solution washed with K_2CO_3 (saturated solution) and water. The organic phase was dried over Na_2SO_4 , filtered and concentrated to dryness. The obtained residue was used without further purification as is described as follows. The obtained residue was dissolved in 15 ml of THF, the solution was cooled to 0°C and 0.601g (0.016 mol) of LiAlH_4 were added in several portions. The reaction mixture was heated under reflux for 4h. After this time, was cooled to 0°C and the excess of hydride was decomposed by the consecutive addition, under stirring, dropwise and cautiously of 0.6 ml of H_2O , 0.6 ml of NaOH (10% solution) and 1.8 ml of H_2O . When the

decomposition was finished, the reaction mixture was filtered and the sludge was washed with THF and CHCl_3 , the organic solutions were combined and concentrated to dryness to obtain 830 mg (75%) of the title product, structure confirmed by ^1H -RMN and ^{13}C -RMN.

- 5 ^{13}C -RMN (CDCl_3): δ 57.0 (CH), 56.5 (CH_2), 47.5 (CH_2), 46.9 (CH_2), 34.3 (CH_3), 26.1 (CH_2), 24.6 (CH), 19.9 (CH_2).

The following Examples of formula (I) were prepared by method (a) from the appropriate carboxylic acid:

- 10 **N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2,3-diphenylpropanamide (Example 25)**
from 2,3-diphenylpropionic acid.
N-(1-Azabicyclo[2.2.2]oct-3-yl)-9H-xanthene-9-carboxamide (Example 28)
N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-9H-xanthene-9-carboxamide (Example 31)
N-[(3S)-1-Azabicyclo[2.2.2]oct-3-yl]-9H-xanthene-9-carboxamide (Example 33)
15 All three from 9H-xanthene-9 carboxylic acid.

Method (b)

Preparation of N-[(3R)-1-azabicyclo[2.2.2]oct-3-yl]-2,2-dithien-2-ylacetamide

- 20 **(Example 17)**
2,2-dithien-2-ylacetic acid was produced by hydrolysis of 2,2-dithien-2-ylacetic acid methyl ester, previously prepared as described in F. Leonard and I. Ehranthal, *J. Am. Chem. Soc.*, (1951), Vol 73, pag 2216. 1.2 g (0.0054 mol) of 2,2-dithien-2-ylacetic acid were dissolved in 25 ml of THF. To this solution were added 0.96 g (0.00594 mol) of 1,1'-
25 carbonyldiimidazole and the mixture was refluxed for an hour. The reaction was monitored by TLC following the formation of the imidazolide. When the reaction was completed 0.75 g (0.00594 mol) of (3R)-aminoquinuclidine were added. The reaction mixture was refluxed for 16 h, cooled, diluted with ether and washed with water. The organic layer was extracted with HCl 2N, the acid solution basified with K_2CO_3 and extracted with CHCl_3 .
30 The organic phase was dried over Na_2SO_4 and the solvent was evaporated to obtain 0.60 g of an oil which was purified by column chromatography (silica gel, CHCl_3 :MeOH: NH_4OH 90:10:1 as eluent). Appropriate fractions were combined and evaporated to give 0.31 g of the title product (17.3%).
mp: 165°C
35 MS $[\text{M}+1]^+$: 333

¹H-NMR (CDCl₃) : δ 1.43 (m, 2H), 1.63 (m, 2H), 1.92 (m, 1H), 2.4 (m, 1H), 2.65-2.85 (m, 4H), 3.31 (m, 1H), 3.97 (m, 1H), 5.37 (s, 1H), 5.98 (d, 1H, NH), 7.0 (m, 4H), 7.28 (m, 2H).

Preparation of N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-9-hydroxy-9H-fluorene-9-carboxamide (Example 44)

5 g (0.022 mol) of 9-hydroxy-9H-fluorene-9-carboxylic acid were dissolved in 50 ml of dry DMF and 4.2 g (0.026 mol) of 1,1'-carbonyldiimidazole were added. The mixture was stirred during 1 h at room temperature. After this time 3.26 g (0.026 mol) of (3R)-aminoquinuclidine and 0.324 g (0.0027 mol) of DMAP (4-(dimethylamino)pyridine) were added. The reaction mixture was stirred at room temperature during 14 hours, then was concentrated in vacuo to eliminate the DMF and the obtained residue was dissolved in AcOEt. The organic solution was washed with NaHCO₃ (saturated solution) and water. The organic layer was separated, dried over Na₂SO₄ and solvent was evaporated. The obtained product was purified by column chromatography (silica gel, CHCl₃: NH₄OH 100:1 → CHCl₃: MeOH:NH₄OH 80:20:1 as eluent). Appropriate fractions were combined and evaporated to yield 304 mg of the title product (4%).

MS [M+1]⁺: 335

¹H-RMN (CDCl₃): δ 7.67 - 7.62 (m, 2H), 7.42 - 7.36 (m, 4H), 7.32 - 7.25 (m, 2H), 6.08 - 5.93 (m, NH, 1H), 3.74 (m, 1H), 3.00 - 2.87 (m, 1H), 2.52 - 2.27 (m, 4H), 2.02 - 1.92 (m, 1H), 1.80 - 1.76 (m, 1H), 1.51 - 1.44 (m, 2H), 1.26 - 1.17 (m, 2H).

Preparation of (2S)-N-[(3R)-1-azabicyclo[2.2.2]oct-3-yl]-2-cyclopentyl-2-hydroxy-2-thien-2-ylacetamide (Example 49)

The title compound was prepared as is described in Example 44 starting from (2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-yl acetic acid. The obtained product was purified by column chromatography (silica gel, CHCl₃: MeOH: NH₄OH 99:1:1 → CHCl₃: MeOH:NH₄OH 97:3:1 as eluent) to give 800 mg of the title product impurified with imidazole. The obtained product was dissolved in CHCl₃ and washed with H₂O (x 3). The organic solution was dried over Na₂SO₄, filtered and concentrated in vacuo to yield 650 mg of the title product (44% from the starting acid). Structure confirmed by ¹H-RMN and MS.

MS [M+1]⁺: 335

¹H-RMN (CDCl₃): δ 7.24 (m, 1H), 7.08 (m, 1H), 6.97 (m, 1H), 6.64 (d, NH, 1H), 3.91 - 3.81 (m, 1H), 3.33 - 3.23 (m, 1H), 2.88 - 2.69 (m, 5H), 2.42 - 2.33 (m, 1H), 1.87 (m, 1H), 1.69 - 1.36 (m, 12H)

(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-yl acetic acid has been prepared as is described in WO/02053564

- 5 The following Examples of formula (I) were prepared according to method (b) from the appropriate carboxylic acid:

**N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-cyclopentyl-2-hydroxy-2-thien-2-ylacetamide
(Example 19)**

- 10 from 2-cyclopentyl-2-hydroxy-2-thien-2-ylacetic acid, which was produced by hydrolysis of 2-cyclopentyl-2-hydroxy-2-thien-2-ylacetic acid methyl ester, compound described in E. Atkinson et al., *J. Med. Chem.*, (1977), Vol 20, n° 12, 1612-1617 and WO02/053564.

**N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-cyclopentyl-2-hydroxy-2-phenylacetamide
(Example 26)**

from 2-cyclopentyl-2-hydroxy-2-phenylacetic acid.

N-(1-Azabicyclo[2.2.2]oct-3-yl)-9H-xanthene-9-carboxamide (Example 28)

from 9H-xanthene-9 carboxylic acid

N-(1-Azabicyclo[2.2.2]oct-3-yl)-9-hydroxy-9H-fluorene-9-carboxamide (Example 41)

- 20 from 9-Hydroxy-9H-fluorene-9-carboxylic acid.

Method (c)

**Preparation of N-[(3R)-1-azabicyclo[2.2.2]oct-3-yl]-2-oxo-2-thien-2-ylacetamide
(Intermediate I-1)**

- 25 Oxalyl chloride (4.5 ml, 0.0516 mol) was added to a solution of 6.68 g (0.0428 mol) of 2-oxo-2-thien-2-ylacetic acid and DMF (several drops) in 100 ml of CHCl₃ (ethanol free) at 0°C. The mixture was stirred and allowed to warm to room temperature. After two hours at this temperature, the solvents were evaporated and the residue was dissolved in CHCl₃
30 and evaporated again. This procedure was repeated two times. The oil obtained was dissolved in CHCl₃ and the solution obtained cooled at 0°C. A solution of (3R)-aminoquinuclidine (5.91g, 0.0468 mol) in 50 ml of CHCl₃ was added. The mixture was stirred and allowed to warm to room temperature. After 18 hours at this temperature, the reaction mixture was washed with an aqueous K₂CO₃ solution. The basic aqueous

solution was extracted again with CHCl_3 . The organic phases were combined, washed with water, dried over Na_2SO_4 and evaporated to give 11.34 g of the title product.

MS $[\text{M}+1]^+$: 265

$^1\text{H-NMR}$ (CDCl_3): δ 1.40-1.85 (m, 4H), 2.0 (m, 1H), 2.60 (m, 1H), 2.70-3.0 (m, 4H), 3.4
5 (m, 1H), 4.0 (m, 1H), 7.20 (m, 1H), 7.50 (d, 1H, NH), 7.85 (m, 1H), 8.40 (m, 1H).

Following an equivalent procedure the following intermediates were prepared:

N-[(3S)-1-Azabicyclo[2.2.2]oct-3-yl]-2-oxo-2-thien-2-ylacetamide (Intermediate I-2)

N-1-Azabicyclo[2.2.2]oct-3-yl]-2-oxo-2-thien-2-ylacetamide (Intermediate I-3)

10

Preparation of N-[(3R)-1-azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2,2-dithien-2-yl-acetamide (Example 4)

A solution of 2-thienylmagnesium bromide was prepared from 2.27 g (0.094 mol) of magnesium and 15.4 g (0.094 mol) of 2-bromothiophene in 150 ml of THF. This solution
15 was added to a solution of 11.34 g (0.043 mol) of N-[(3R)-1-azabicyclo[2.2.2]oct-3-yl]-2-oxo-2-thien-2-ylacetamide (Intermediate I-1) in 120 ml of THF. The mixture was stirred at room temperature for one hour, refluxed for one hour, cooled and treated with a saturated solution of ammonium chloride. The solution obtained was basified with a K_2CO_3 solution and extracted with CH_2Cl_2 . The organic layer was washed with water and brine, dried over
20 Na_2SO_4 and treated with activated carbon. The solution obtained was filtered through Celite and solvents were evaporated to give a solid that was treated with ether and filtered to give 8.77 (58.5%) of the title product.

m. p.: 169°C

MS $[\text{M}+1]^+$: 349

25 $^1\text{H-NMR}$ ($\text{DMSO}-d_6$): δ 1.30 (m, 1H), 1.42-1.75 (m, 3H), 1.77 (m, 1H), 2.50-2.75 (m, 4H), 2.83 (m, 1H), 3.05 (m, 1H), 3.74 (m, 1H), 6.95 (m, 2H), 7.03 (m, 1H), 7.09 (m, 1H), 7.43 (m, 2H), 7.54 (s, 1H, OH), 7.97 (d, 1H, NH).

The following Examples of formula (I) were prepared according to method (c) using N-
30 [(3R)-1-azabicyclo[2.2.2]oct-3-yl]-2-oxo-2-thien-2-ylacetamide (Intermediate I-1) and the corresponding Grignard reagent:

N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2-thien-2-ylpent-4-enamide (Example 20).

(2*)-N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2-thien-2-ylbutanamide
35 **(diastereomer 1, Example 21) and**

**(2*)-N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2-thien-2-ylbutanamide
(diastereomer 2, Example 22)**

the two diastereomers were separated by column chromatography (silica gel,
CHCl₃/MeOH/NH₄OH 90:10:1 as eluent). Diastereomer 1: first diastereomer obtained,

5 Diastereomer 2: second distereomer obtained.

**(2*)-N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2-thien-2-ylbut-3-enamide
(diastereomer 1, Example 23) and**

**(2*)-N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2-thien-2-ylbut-3-enamide
(diastereomer 2, Example 24)**

10 the two diastereomers were separated by column chromatography (silica gel,
CHCl₃/MeOH/NH₄OH 90:10:1 as eluent). Diastereomer 1: first diastereomer obtained,
Diastereomer 2: second distereomer obtained.

(*) Configuration not assigned.

15

The following compounds of formula (I) were also prepared according to method (c)
**N-[(3S)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2,2-dithien-2-ylacetamide (Example
15)**

prepared from Intermediate I-2 and 2-thienyllithium (commercially available).

20 **N-(1-Azabicyclo[2.2.2]oct-3-yl)-2-hydroxy-2,2-dithien-2-ylacetamide (Example 1)**
prepared from Intermediate I-3 and 2-thienyllithium.

**Preparation of N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-(5-bromothien-2-yl)-2-(4-fluoro-
3-methylphenyl)-2-hydroxyacetamide (Example 48)**

25 Oxalyl chloride (1.16 ml, 0.013 mol) was added to a suspension of 2-(5-bromothien-2-yl)-
2-oxoacetic acid (2.6 g, 0.011 mol) and two drops of DMF in 30 ml of CHCl₃ (ethanol free)
cooled to 0°C. The mixture was allowed to warm to room temperature under stirring.
After 1.5 hours the reaction mixture was concentrated to dryness in vacuo and the
obtained residue was dissolved in CHCl₃ (15 ml) and concentrated again. This procedure
30 was repeated two times. The obtained residue was dissolved in CHCl₃ (30 ml), the
solution was cooled to 0°C and 1.67 g (0.013 mol) of (3R)-aminoquinuclidine were
added. The mixture was allowed to warm to room temperature and continued stirring
during 3 hours. After this time the reaction mixture was treated with K₂CO₃ (saturated
solution) and the aqueous layer was extracted with CH₂Cl₂. The organic layers were
35 combined, washed with water, dried over Na₂SO₄, filtered and evaporated. The obtained

product (N-[(3R)-1-azabicyclo[2.2.2]oct-3-yl]-2-(5-bromothien-2-yl)-2-oxoacetamide) was used without further purification as is described as follows.

The obtained product was dissolved in dry THF (45 ml). The solution obtained was cooled to -80°C and 14.4 ml of a 1M solution in THF of 4-fluoro-3-methylphenylmagnesium bromide (0.014 mol) were added. The mixture was stirred during 3 hours at low temperature. After this time, the reaction mixture was treated with NH_4Cl (saturated solution) and extracted with AcOEt. The organic layer was dried over Na_2SO_4 , filtered and solvents were evaporated. The obtained product was purified by column chromatography (silica gel, CHCl_3 : MeOH: NH_4OH 99:1:1 \rightarrow CHCl_3 : MeOH: NH_4OH 96:4:1) to yield 2.1g of the title compound (42% from the starting acid) as a mixture of diastereomers.

MS $[\text{M}+1]^+$: 453, 455

^1H -RMN (CDCl_3): (mixture of diastereomers) δ 7.35 - 7.20 (m, 2H), 7.08 - 6.89 (m, 3H), 6.80 (dd, 1H), 3.93 - 3.84 (m, 1H), 3.18 - 3.07 (m, 1H), 2.69 - 2.58 (m, 4H), 2.33 - 2.22 (m, 4H), 1.92 - 1.87 (m, 1H), 1.63 - 1.42 (m, 4H)

2-(5-bromothien-2-yl)-2-oxoacetic acid was obtained by hydrolysis of ethyl 2-(5-bromothien-2-yl)-2-oxoacetate by a standard method (EtOH , NaOH 2N, 60°C , 1.5 h).

Preparation of N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-fur-2-yl-2-hydroxypent-3-ynamide (Example 45)

To a suspension cooled to 0°C of 1.72 g (0.012 mol) of 2-fur-2-yl-2-oxoacetic acid and two drops of DMF in 40 ml of CHCl_3 (ethanol free), 1.17 ml (0.013 mol) of oxalyl chloride were added. The resulting mixture was allowed to warm to room temperature under stirring and continued stirring during 1 hour. After this time, the mixture was concentrated to dryness in vacuo and the obtained residue was dissolved in CHCl_3 (20 ml) and concentrated again. This procedure was repeated two times. The obtained product was dissolved in CHCl_3 (50 ml), the solution was cooled to 0°C and 1.7 g (0.013 mol) of (3R)-aminoquinuclidine were added. The mixture was allowed to warm to room temperature and continued stirring during 16 hours. After this time the reaction mixture was treated with K_2CO_3 (saturated solution) and the aqueous layer was extracted with CH_2Cl_2 . The organic layers were combined, washed with water, dried over Na_2SO_4 , filtered and evaporated. The obtained product (N-[(3R)-1-azabicyclo[2.2.2]oct-3-yl]-2-(2-furyl)-2-oxoacetamide) was used without further purification as is described as follows. This product was dissolved in dry THF (50 ml). The solution obtained was cooled to -80°C and 28 ml of a 0.5 M solution in THF of 1-propynylmagnesium bromide (0.014 mol) were

added. The mixture was allowed to warm to room temperature during 3 hours. After this time, the reaction mixture was treated with NH_4Cl (saturated solution) and extracted with AcOEt and CH_2Cl_2 . The organic layers were combined, dried over Na_2SO_4 , filtered and solvents were evaporated. The obtained product was purified by column chromatography (silica gel, CHCl_3 : MeOH : NH_4OH 97:3:1 \rightarrow CHCl_3 : MeOH : NH_4OH 94:6:1 as eluent) to yield 1.38 g of the title compound (39% from the starting acid) as a mixture of diastereomers.

MS $[\text{M}+1]^+$: 289

^1H -RMN (CDCl_3): (mixture of diastereomers) δ 7.36 - 7.35 (m, 1H), 6.87 - 6.76 (m, NH, 1H), 6.54 - 6.52 (m, 1H), 6.34 - 6.32 (m, 1H), 3.92 (m, 1H), 3.29 - 3.17 (m, 1H), 2.78 - 2.64 (m, 4H), 2.47 - 2.35 (m, 1H), 2.00 - 1.90 (m, 1H), 1.92 (s, 3H), 1.70 - 1.39 (m, 4H).

Preparation of N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-fur-2-yl-2-hydroxy-4-(4-methoxyphenyl)butanamide (Example 46)

This product was prepared according to the procedure described in Example 45, starting from 2 g (0.014 mol) of 2-fur-2-yl-2-oxoacetic acid, and using 4-methoxyphenethylmagnesium chloride (0.021 mol, 43 ml of a 0.5 M solution in THF) as Grignard reagent. The obtained product was purified by column chromatography (silica gel, CHCl_3 : MeOH : NH_4OH 99:1:1 \rightarrow CHCl_3 : MeOH : NH_4OH 97:3:1 as eluent) to yield 2.4 g of the title compound (44% from the starting acid) as a mixture of diastereomers.

MS $[\text{M}+1]^+$: 385

^1H -RMN (CDCl_3): (mixture of diastereomers) δ 7.39 - 7.38 (m, 1H), 7.09 (d, 2H), 6.82 (d, 2H), 6.70 - 6.67 (m, NH, 1H), 6.41 - 6.35 (m, 2H), 3.92 (m, 1H), 3.78 (s, 3H), 3.33 - 3.24 (m, 1H), 2.82 - 2.19 (m, 9H), 1.96 - 1.87 (m, 1H), 1.67 - 1.40 (m, 4H)

Method (d)

Preparation of (3R)-1-(3-phenoxypropyl)-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane bromide (Example 32)

0.3 g (0.00089 mol) of N-[(3R)-1-azabicyclo[2.2.2]oct-3-yl]-9H-xanthene-9-carboxamide (Example 31) were dissolved in 6 ml of THF. To this solution 0.21 ml (0.287g, 0.001335 mol) of (3-bromopropoxy)benzene were added. The mixture was refluxed during 4 hours and then stirred 17 h at room temperature. The reaction mixture was filtered and the solid obtained washed with ether several times and dried. 0.48 g (98%) of the title product were obtained.

m.p.: 278.5-279.2°C

MS [M-Br]⁺: 469

¹H-NMR (DMSO-d₆): δ 1.80-2.0 (m, 3H), 2.05-2.30 (m, 4H), 3.15-3.25 (m, 1H), 3.30-3.65 (m, 6H), 3.82 (m, 1H), 4.05 (t, 2H), 4.10 (m, 1H), 5.05 (s, 1H), 6.90-7.0 (m, 3H), 7.05-7.20 (m, 4H), 7.25-7.42 (m, 6H), 9.01 (d, 1H, NH).

Preparation of 3-[[[9-hydroxy-9H-fluoren-9-yl]carbonyl]amino]-1-methyl-1-azoniabicyclo[2.2.2]octane bromide (Example 42)

0.245 g (0.00073 mol) of N-(1-azabicyclo[2.2.2]oct-3-yl)-9-hydroxy-9H-fluorene-9-carboxamide (Example 41) were dissolved in 4 ml of acetonitrile and 6 ml of CHCl₃. To this solution 5.46 ml of a 1 M solution of methyl bromide in acetonitrile were added and the mixture was stirred for 96 h at room temperature under N₂ atmosphere. After this time the solvents were evaporated. The residue was coevaporated with ether three times. Ether was added to the residue and the mixture stirred. The solid obtained was filtered and washed several times with ether. The yield was 0.26 g (83.8%) of the title compound.

m.p.: 197.5-203.6°C

MS [M-Br]⁺: 349

¹H-NMR (DMSO-d₆): δ 1.80-2.0 (m, 3H), 2.10-2.25 (m, 2H), 2.96 (s, 3H), 3.35-3.70 (m, 5H), 3.78 (m, 1H), 4.15 (m, 1H), 6.82 (s, 1H, OH), 7.30 (m, 2H), 7.43 (m, 4H), 7.79 (m, 2H), 8.81 (d, 1H, NH).

Method (e)

Preparation of (3S)-1-cyclohexylmethyl-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane trifluoroacetate (Example 36)

28.15 mg (0.0842 mmol) of N-[(3S)-1-azabicyclo[2.2.2]oct-3-yl]-9H-xanthene-9-carboxamide (Example 33) were dissolved in 0.5 ml of DMSO. A solution of 74.37 mg (0.421 mmol) of (bromomethyl)cyclohexane in 0.5 ml of DMSO was added and the mixture stirred at room temperature overnight. The mixture was purified by solid phase extraction with a cation exchange Mega Bond Elut cartridge, previously conditioned at pH=7.5 with 0.1 M NaH₂PO₄ buffer. The reaction mixture was applied to the cartridge and washed first with 2 ml of DMSO and then three times with 5 ml of acetonitrile, rinsing away all starting materials. The ammonium derivative was eluted with 5 ml of 0.03 M TFA solution in CH₃CN:CHCl₃ (2:1). This solution was neutralized with 300 mg of poly(4-

vinylpyridine), filtered and evaporated to dryness. The yield was 12.4 mg of the title product. (27.0%).

[M-CF₃COO]⁺: 431

5 Preparation of (3R)-3-(2-hydroxy-2,2-dithien-2-ylacetyl-amino)-1-[3-(3-hydroxyphenoxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate (Example 12)

The title compound was prepared as described for Example 36 from 30 mg (0.0861 mmol) of N-[(3R)-1-azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2,2-dithien-2-ylacetamide (Example 4)
10 (dissolved in 0.5 ml of DMSO) and 78 mg (0.418 mmol) of 3-(3-chloropropoxy)phenol (dissolved in 0.5 ml of DMSO). The yield was 12.7 mg of the title product. (24%).
[M-CF₃COO]⁺: 499

Also included within the scope of the present invention are pharmaceutical compositions
15 which comprise, as the active ingredient, at least one quinuclidine amide derivative of formulae (I) or (II) in association with a pharmaceutically acceptable carrier or diluent. Preferably the composition is made up in a form suitable for oral administration.

The pharmaceutically acceptable carrier or diluents which are mixed with the active
20 compound or compounds, to form the composition of this invention are well-known per se and the actual excipients used depend inter alia on the intended method of administration of the composition.

Compositions of this invention are preferably adapted for oral administration. In this case,
25 the composition for oral administration may take the form of tablets, film-coated tablets, liquid inhalant, powder inhalant and inhalation aerosol; all containing one or more compounds of the invention; such preparations may be made by methods well-known in the art.

30 The diluents which may be used in the preparations of the compositions include those liquid and solid diluents which are compatible with the active ingredient, together with colouring or flavouring agents, if desired. Tablets or film-coated tablets may conveniently contain between 1 and 500 mg, preferably from 5 to 300 mg of active ingredient. The inhalant compositions may contain between 1 µg and 1,000 µg, preferably from 10 µg to
35 800 µg of active ingredient. In human therapy, the dose of the compound of formulae (I) or

(II) depend on the desired effect and duration of treatment; adult doses are generally between 3 mg and 300 mg per day as tablets and 10 µg and 800 µg per day as inhalant composition.

5 **Pharmacological Action**

The results on human muscarinic receptors binding and in the test on bronchospasm in guinea pig, were obtained as described below.

10 **Human muscarinic receptor studies.**

The binding of [3H]-NMS to human muscarinic receptors was performed according to Waelbroeck et al (1990), Mol. Pharmacol., 38: 267-273. Assays were carried out at 25°C. Membrane preparations from stably transfected chinese hamster ovary-K1 cells (CHO)

15 expressing the genes for the human M3 muscarinic receptors were used.

For determination of IC₅₀, membrane preparations were suspended in DPBS to a final concentration of 89 µg/ml for the M3 subtype. The membrane suspension was incubated with the tritiated compound for 60 min. After incubation the membrane fraction was

20 separated by filtration and the bound radioactivity determined. Non-specific binding was determined by addition of 10⁻⁴ M atropine. At least six concentrations were assayed in duplicate to generate individual displacement curves.

Our results show that the compounds of the present invention have high affinities for M3

25 muscarinic receptors, preferably human muscarinic receptors. Thus, the IC₅₀ of the preferred compounds of the invention is lower than 100 nM. Most preferred compounds, such as the compounds of examples 1, 3, 4, 8, 10, 11, 14, 19, 30 and 32 described below, have an IC₅₀ lower than 60 nM.

30 **Test on bronchospasm in guinea pig**

The studies were performed according to H. Konzett and F. Rössler (1940), Arch. Exp. Path. Pharmacol. 195: 71-74. Aqueous solutions of the agents to be tested were nebulized and inhaled by anaesthetized ventilated male guinea pigs (Dunkin-Hartley).

35 Bronchial response to intravenous acetylcholine challenge was determined before and

after drug administration and the changes in pulmonary resistance at several time-points were expressed as percent of inhibition of bronchospasm.

The compounds of the present invention inhibited the bronchospasm response to
5 acetylcholine with high potency and a long duration of action.

From the above described results one of ordinary skill in the art can readily understand that the compounds of the present invention have excellent M3 antimuscarinic activity and thus are useful for the treatment of diseases in which the M3 muscarinic receptor is
10 implicated, including respiratory disorders such as chronic obstructive pulmonary disease (COPD), bronchitis, bronchial hyperreactivity, asthma, cough and rhinitis; urological disorders such as urinary incontinence, pollakiuria, neurogenic or unstable bladder, cystospasm and chronic cystitis; gastrointestinal disorders such as irritable bowel syndrome, spastic colitis, diverticulitis and peptic ulceration; and cardiovascular disorders
15 such as vagally induced sinus bradycardia.

The present invention further provides a compound of formulae (I) or (II) or a pharmaceutically acceptable composition comprising a compound of formulae (I) or (II) for use in a method of treatment of the human or animal body by therapy, in particular for the
20 treatment of respiratory, urological or gastrointestinal disease or disorder.

The present invention further provides the use of a compound of formulae (I) or (II) or a pharmaceutically acceptable composition comprising a compound of formulae (I) or (II) for the manufacture of a medicament for the treatment of a respiratory, urological or
25 gastrointestinal disease or disorder.

Further, the compounds of formulae (I) or (II) and pharmaceutical compositions comprising a compound of formulae (I) or (II) can be used in a method of treating a respiratory, urological or gastrointestinal disease or disorder, which method comprises
30 administering to a human or animal patient in need of such treatment an effective, non-toxic, amount of a compound of formulae (I) or (II) or a pharmaceutical composition comprising a compound of formulae (I) or (II).

Further, the compounds of formulae (I) or (II) and pharmaceutical compositions
35 comprising a compound of formulae (I) or (II) can be used in combination with other drugs

effective in the treatment of these diseases. For example with β_2 agonists, steroids, antiallergic drugs, phosphodiesterase IV inhibitors and/or leukotriene D4 (LTD4) inhibitors, for simultaneous, separate or sequential use in the treatment of a respiratory disease.

- 5 The present invention will be further illustrated by the following examples. The examples are given by way of illustration only and are not to be construed as a limiting.

Example 1

N-(1-Azabicyclo[2.2.2]oct-3-yl)-2-hydroxy-2,2-dithien-2-ylacetamide

- 10 The title compound was synthesised according to method c. The yield was 0.67 g , 15.2%.
m.p.: 185°C.

MS [M+1]⁺: 349.

- ¹H-NMR (DMSO-d₆): δ 1.30 (m, 1H), 1.40-1.75 (m, 3H), 1.77 (m, 1H), 2.50-2.75 (m, 4H),
2.83 (m, 1H), 3.05 (m, 1H), 3.74 (m, 1H), 6.94 (m, 2H), 7.04 (m, 1H), 7.10 (m, 1H), 7.43
15 (m, 2H), 7.53 (s, 1H, OH), 7.95 (d, 1H, NH).

Example 2

3-(2-Hydroxy-2,2-dithien-2-ylacetyl-amino)-1-methyl-1-azoniabicyclo[2.2.2]octane bromide

- 20 The title compound was synthesised according to methods c and d. The yield of the final step was 0.16 g, 68%.

MS [M-Br]⁺: 363.

- ¹H-NMR (DMSO-d₆): δ 1.76 (m, 1H), 1.85-2.08 (m, 3H), 2.14 (m, 1H), 2.92 (s, 3H), 3.30-
3.60 (m, 5H), 3.75 (m, 1H), 4.20 (s, 1H), 7.0 (m, 2H), 7.06 (m, 1H), 7.12 (m, 1H), 7.46 (m,
25 2H), 7.65 (s, 1H, OH), 8.64 (d, 1H, NH).

Example 3

3-(2-Hydroxy-2,2-dithien-2-ylacetyl-amino)-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane bromide

- 30 The title compound was synthesised according to methods c and d. The yield of the final step was 0.24 g, 82.7%.

m.p.: 180.6-188.3°C.

MS [M-Br]⁺: 483.

¹H-NMR (DMSO-d₆): δ 1.78 (m, 1H), 1.90-2.25 (m, 6H), 3.30-3.65 (m, 7H), 3.67-3.80 (m, 1H), 4.05 (t, 2H), 4.24 (m, 1H), 6.94-7.0 (m, 5H), 7.07 (m, 1H), 7.13 (m, 1H), 7.31 (m, 2H), 7.47 (m, 2H), 7.66 (s, 1H, OH), 8.74 (d, 1H, NH).

5 Example 4

N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2,2-dithien-2-ylacetamide

The title compound was synthesised according to method c. The yield was 8.77 g, 58.5%.
m.p.: 169°C.

MS [M+1]⁺: 349.

- 10 ¹H-NMR (DMSO-d₆): δ 1.30 (m, 1H), 1.42-1.75 (m, 3H), 1.77 (m, 1H), 2.50-2.75 (m, 4H), 2.83 (m, 1H), 3.05 (m, 1H), 3.74 (m, 1H), 6.95 (m, 2H), 7.03 (m, 1H), 7.09 (m, 1H), 7.43 (m, 2H), 7.54 (s, 1H, OH), 7.97 (d, 1H, NH).

Example 5

- 15 **(3R)-3-(2-Hydroxy-2,2-dithien-2-ylacetyl-amino)-1-methyl-1-azoniabicyclo[2.2.2]octane trifluoroacetate**

The title compound was synthesised according to methods c and e. The yield of the final step was 0.0124 g, 30.2%.

MS [M-CF₃COO]⁺: 363.

20

Example 6

(3R)-1-Allyl-3-(2-hydroxy-2,2-dithien-2-ylacetyl-amino)-1-azoniabicyclo[2.2.2]octane trifluoroacetate

- 25 The title compound was synthesised according to methods c and e. The yield of the final step was 0.010 g, 23.1%.

MS [M-CF₃COO]⁺: 389.

Example 7

- 30 **(3R)-1-Heptyl-3-(2-hydroxy-2,2-dithien-2-ylacetyl-amino)-1-azoniabicyclo[2.2.2]octane trifluoroacetate**

The title compound was synthesised according to methods c and e. The yield of the final step was 0.0118 g, 24.4%.

MS [M-CF₃COO]⁺: 447.

Example 8**(3R)-3-(2-Hydroxy-2,2-dithien-2-ylacetyl-amino)-1-(3-phenylpropyl)-1-azoniabicyclo[2.2.2]octane bromide**

The title compound was synthesised according to methods c and d. The yield of the final
5 step was 0.49 g, 78%.

m.p.: 117.3-118.9°C.

MS [M-Br]⁺: 467

¹H-NMR (DMSO-d₆): δ 1.75 (m, 1H), 1.80-2.05 (m, 5H), 2.09 (m, 1H), 2.60 (m, 2H), 3.16
(m, 2H), 3.25-3.60 (m, 5H), 3.68 (m, 1H), 4.21 (m, 1H), 6.96 (m, 2H), 7.04 (m, 1H), 7.11
10 (m, 1H), 7.20-7.40 (m, 5H), 7.45 (m, 2H), 7.62 (s, 1H, OH), 8.70 (d, 1H, NH).

Example 9**(3R)-3-(2-Hydroxy-2,2-dithien-2-ylacetyl-amino)-1-((E)-3-phenylallyl)-1-azoniabicyclo[2.2.2]octane trifluoroacetate**

15 The title compound was synthesised according to methods c and e. The yield of the final
step was 0.0032 g, 6.4%.

[M-CF₃COO]⁺: 465.

Example 10**20 (3R)-3-(2-Hydroxy-2,2-dithien-2-ylacetyl-amino)-1-(2-phenoxyethyl)-1-azoniabicyclo[2.2.2]octane bromide**

The title compound was synthesised according to methods c and d. The yield of the final
step was 0.470 g, 74%.

m.p.: 112.6-113.9°C.

25 MS [M-Br]⁺: 469

¹H-NMR (DMSO-d₆): δ 1.77 (m, 1H), 1.90-2.15 (m, 4H), 3.40-3.80 (m, 7H), 3.86 (m, 1H),
4.24 (m, 1H), 4.43 (m, 2H), 6.95-7.0 (m, 5H), 7.04 (m, 1H), 7.11 (m, 1H), 7.35 (m, 2H),
7.46 (m, 2H), 7.66 (s, 1H, OH), 8.72 (d, 1H, NH).

30 Example 11**(3R)-3-(2-Hydroxy-2,2-dithien-2-ylacetyl-amino)-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane bromide**

The title compound was synthesised according to methods c and d. The yield of the final
step was 0.65 g, 80%.

35 m.p.: 182°C.

MS [M-Br]⁺: 483

¹H-NMR (DMSO-d₆): δ 1.76 (m, 1H), 1.85-2.25 (m, 6H), 3.25-3.65 (m, 7H), 3.65-3.82 (m, 1H), 4.05 (t, 2H), 4.23 (m, 1H), 6.90-7.0 (m, 5H), 7.04 (m, 1H), 7.11 (m, 1H), 7.31 (m, 2H), 7.45 (m, 2H), 7.65 (s, 1H, OH), 8.73 (d, 1H, NH).

5

Example 12

(3R)-3-(2-Hydroxy-2,2-dithien-2-ylacetyl-amino)-1-[3-(3-hydroxyphenoxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was synthesised according to methods c and e. The yield of the final
10 step was 0.0127 g, 24.0%.

[M-CF₃COO]⁺: 499.

Example 13

**(3R)-1-(2-Benzoyloxyethyl)-3-(2-hydroxy-2,2-dithien-2-ylacetyl-amino)-1-
15 azoniabicyclo[2.2.2]octane trifluoroacetate**

The title compound was synthesised according to methods c and e. The yield of the final
step was 0.0146 g, 28.4%.

[M-CF₃COO]⁺: 483.

20 Example 14

**(3R)-3-(2-Hydroxy-2,2-dithien-2-ylacetyl-amino)-1-(3-thien-2-ylpropyl)-1-
azoniabicyclo[2.2.2]octane bromide**

The title compound was synthesised according to methods c and d. The yield of the final
step was 0.49 g, 77%.

25 m.p.: 111.1-113.2°C.

MS [M-Br]⁺: 473

¹H-NMR (DMSO-d₆): δ 1.75 (m, 1H), 1.85-2.15 (m, 6H), 2.84 (t, 2H), 3.17 (m, 2H), 3.25-
3.45 (m, 3H), 3.45-3.62 (m, 2H), 3.70 (m, 1H), 4.21 (m, 1H), 6.90-7.0 (m, 4H), 7.05 (m,
1H), 7.11 (m, 1H), 7.38 (m, 1H), 7.45 (m, 2H), 7.65 (s, 1H, OH), 8.71 (d, 1H, NH).

30

Example 15

N-[(3S)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2,2-dithien-2-ylacetamide

The title compound was synthesised according to method c. The yield was 0.44 g, 7.67%.

MS [M+1]⁺: 349.

¹H-NMR (CDCl₃): δ 1.40-1.70 (m, 4H), 1.98 (m, 1H), 2.40-2.47 (m, 1H), 2.60-2.85 (m, 4H), 3.24 (m, 1H), 3.97 (m, 1H), 6.91 (d, 1H), 7.01 (m, 2H), 7.15 (m, 2H), 7.33 (m, 2H).

Example 16

5 **(3S)-3-(2-Hydroxy-2-dithien-2-ylacetyl-amino)-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane bromide**

The title compound was synthesised according to methods c and d. The yield of the final step was 0.36 g, 70.6%.

m.p.: 172.8-173.9°C.

10 MS [M-Br]⁺: 483.

¹H-NMR (DMSO-d₆): δ 1.78 (m, 1H), 1.85-2.25 (m, 6H), 3.25-3.65 (m, 7H), 3.70-3.82 (m, 1H), 4.04 (t, 2H), 4.24 (m, 1H), 6.90-7.0 (m, 5H), 7.06 (m, 1H), 7.12 (m, 1H), 7.31 (m, 2H), 7.45 (m, 2H), 7.65 (s, 1H, OH), 8.74 (d, 1H, NH).

15 Example 17

N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2,2-dithien-2-ylacetamide

The title compound was synthesised according to method b. The yield was 0.31 g, 17.3%.

m.p.: 165°C.

MS [M+1]⁺: 333.

20 ¹H-NMR (CDCl₃): δ 1.43 (m, 2H), 1.63 (m, 2H), 1.92 (m, 1H), 2.4 (m, 1H), 2.65-2.85 (m, 4H), 3.31 (m, 1H), 3.97 (m, 1H), 5.37 (s, 1H), 5.98 (d, 1H, NH), 7.0 (m, 4H), 7.28 (m, 2H).

Example 18

(3R)-3-(2,2-Dithien-2-ylacetyl-amino)-1-(3-phenoxypropyl)-1-

25 **azoniabicyclo[2.2.2]octane bromide**

The title compound was synthesised according to methods b and d. The yield of the final step was 0.21 g, 79.8%.

m.p.: 135.6-137.1°C.

MS [M-Br]⁺: 467.

30 ¹H-NMR (DMSO-d₆): δ 1.83 (m, 1H), 1.94 (m, 2H), 2.0-2.25 (m, 4H), 3.19 (m, 1H), 3.30-3.55 (m, 6H), 3.86 (m, 1H), 4.03 (t, 2H), 4.18 (m, 1H), 5.65 (s, 1H), 6.93-6.98 (m, 5H), 7.01-7.06 (m, 2H), 7.31 (m, 2H), 7.42 (m, 2H), 9.10 (d, 1H, NH).

Example 19**N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-cyclopentyl-2-hydroxy-2-thien-2-ylacetamide**

The title compound was synthesised as a mixture of diastereomers according to method b. The yield was 0.12 g, 27.9%.

5 MS [M+1]⁺: 335.

¹H-NMR (DMSO-d₆): (mixture of diastereomers 50:50) δ 1.20-1.80 (m, 13H), 2.50-2.90 (m, 6H), 3.04 (m, 1H), 3.69 (m, 1H), 5.98 & 6.01 (s, 1H, OH), 6.93 (m, 1H), 7.08 (m, 1H), 7.35 (m, 1H), 7.59 (m, 1H, NH).

10 **Example 20**

N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2-thien-2-ylpent-4-enamide

The title compound was synthesised as a mixture of diastereomers according to method c. The yield was 1.92 g, 82.4%.

m.p.: 54.4-58.3°C

15 MS [M+1]⁺: 307.

¹H-NMR (DMSO-d₆): (mixture of diastereomers) δ 1.25 (m, 1H), 1.40-1.75 (m, 4H), 2.45-2.70 (m, 5H), 2.70-3.10 (m, 3H), 3.65 (m, 1H), 5.0-5.10 (m, 2H), 5.60-5.80 (m, 1H), 6.50 & 6.52 (s, 1H, OH), 6.95 (m, 1H), 7.05 (m, 1H), 7.38 (m, 1H), 7.60 (d, 1H, NH).

20 **Example 21**

(2*)-N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2-thien-2-ylbutanamide

(diastereomer 1, 70:30)

The title compound was synthesised according to method c. The yield was 0.19 g, 34% (based on single isomer).

25 m.p.: 139.0-140.7°C.

MS [M+1]⁺: 295.

¹H-NMR (DMSO-d₆): (diastereomer 1, 70:30) δ 0.80 (t, 3H), 1.27 (m, 1H), 1.40-1.80 (m, 4H), 1.83 (m, 1H), 2.16 (m, 1H), 2.45-2.70 (m, 4H), 2.79 (m, 1H), 2.90-3.10 (m, 1H), 3.67 (m, 1H), 6.28 & 6.31 (s, 1H, OH), 6.94 (m, 1H), 7.03 (m, 1H), 7.36 (m, 1H), 7.62 (d, 1H, NH).

30

(*): Configuration not assigned.

Example 22**(2*)-N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2-thien-2-ylbutanamide**

35 (diastereomer 2, 27:73)

The title compound was synthesised according to method c. The yield was 0.42 g , 75% (based on single isomer).

m.p.: 68.9-70.2°C.

MS [M+1]⁺: 295.

- 5 ¹H-NMR (DMSO-d₆): (diastereomer 2, 27:73) δ 0.82 (t, 3H), 1.26 (m, 1H), 1.40-1.80 (m, 4H), 1.86 (m, 1H), 2.17 (m, 1H), 2.45-2.70 (m, 4H), 2.78 (m, 1H), 2.90-3.10 (m, 1H), 3.67 (m, 1H), 6.29 & 6.32 (s, 1H, OH), 6.93 (m, 1H), 7.03 (m, 1H), 7.36 (m, 1H), 7.62 (d, 1H, NH).

(*): Configuration not assigned.

10

Example 23

(2*)-N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2-thien-2-ylbut-3-enamide

(diastereomer 1)

The title compound was synthesised according to method c. The yield was 0.21 g, 18.9 %

- 15 (based on single isomer).

m.p.: 171.7-173.2°C.

MS [M+1]⁺: 293.

- ¹H-NMR (DMSO-d₆): (diastereomer 1) δ 1.27 (m, 1H), 1.40-1.75 (m, 4H), 2.50-2.70 (m, 4H), 2.77 (m, 1H), 3.0 (m, 1H), 3.66 (m, 1H), 5.18 (d, 1H), 5.38 (d, 1H), 6.41 (dd, 1H),
20 6.71 (s, 1H, OH), 6.95 (m, 1H), 7.02 (m, 1H), 7.41 (m, 1H), 7.70 (d, 1H, NH).

(*): Configuration not assigned.

Example 24

(2*)-N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2-thien-2-ylbut-3-enamide

- 25 (diastereomer 2)

The title compound was synthesised according to method c. The yield was 0.21 g, 18.9 % (based on single isomer)

m.p.: 53.8-55.0°C.

MS [M+1]⁺: 293.

- 30 ¹H-NMR (DMSO-d₆): (diastereomer 2) δ 1.27 (m, 1H), 1.40-1.75 (m, 4H), 2.50-2.70 (m, 4H), 2.81 (m, 1H), 3.03 (m, 1H), 3.69 (m, 1H), 5.21 (d, 1H), 5.40 (d, 1H), 6.45 (dd, 1H), 6.74 (s, 1H, OH), 6.96 (m, 1H), 7.01 (m, 1H), 7.41 (m, 1H), 7.72 (d, 1H, NH).

(*): Configuration not assigned.

Example 25**N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2,3-diphenylpropanamide**

The title compound was synthesised as a mixture of diastereomers according to method a. The yield was 1.21 g, 82.3 %.

5 m.p.: 142°C.

MS [M+1]⁺: 335.

¹H-NMR (DMSO-d₆): (mixture of diastereomers) δ 0.95-1.30 (m, 2H), 1.35-1.50 (m, 2H), 1.52 (m, 1H), 2.14-2.27 (m, 1H), 2.50-2.65 (m, 4H), 2.81-2.96 (m, 2H), 3.24-3.32 (m, 1H), 3.56 (m, 1H), 3.78-3.89 (m, 1H), 7.10-7.45 (m, 10H), 8.13 (m, 1H, NH).

10

Example 26**N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-cyclopentyl-2-hydroxy-2-phenylacetamide**

The title compound was synthesised as a mixture of diastereomers according to method b. The yield was 0.25 g, 28.4 %.

15 m.p.: 69.8-73.3°C.

MS [M+1]⁺: 329.

¹H-NMR (DMSO-d₆): (mixture of diastereomers) δ 1.25 (m, 3H), 1.35-1.75 (m, 10H), 2.40-2.70 (m, 4H), 2.76 (m, 1H), 2.90-3.10 (m, 2H), 3.62 (m, 1H), 5.60 & 5.62 (s, 1H, OH), 7.20-7.23 (m, 1H), 7.27-7.33 (m, 2H), 7.52-7.61 (m, 3H).

20

Example 27**N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-phenylhexanamide**

The title compound was synthesised as a mixture of diastereomers according to method a. The yield was 1.59 g, 63.1 %.

25 m.p.: 136°C.

MS [M+1]⁺: 301.

¹H-NMR (DMSO-d₆): (mixture of diastereomers 50:50) δ 0.84 (t, 3H), 1.10-1.60 (m, 9H), 1.70 (m, 1H), 1.90 (m, 1H), 2.25-2.50 (m, 1H), 2.50-2.80 (m, 4H), 2.94 & 3.05 (m, 1H), 3.50 (m, 1H), 3.65 (m, 1H), 7.15-7.40 (m, 5H), 8.04 (m, 1H, NH).

30

Example 28**N-(1-Azabicyclo[2.2.2]oct-3-yl)-9H-xanthene-9-carboxamide**

The title compound was synthesised according to method b. The yield was 0.28g, 19 %.

m.p.: 251°C.

35 MS [M+1]⁺: 335.

¹H-NMR (DMSO-d₆): δ 1.25-1.60 (m, 3H), 1.67 (m, 1H), 1.82 (m, 1H), 2.40-2.50 (m, 1H), 2.55-2.75 (m, 3H), 2.82 (m, 1H), 3.06 (m, 1H), 3.62 (m, 1H), 5.0 (s, 1H), 7.05-7.15 (m, 4H), 7.25-7.35 (m, 4H), 8.51 (d, 1H, NH).

The title compound was also synthesised according to method a. The yield was 2.11 g, 52.6 %.

Example 29

1-Methyl-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane bromide

The title compound was synthesised according to methods a and d. The yield of the final step was 0.25 g, 77.6%.

m.p.: 318°C.

MS [M-Br]⁺: 349.

¹H-NMR (DMSO-d₆): δ 1.80-2.0 (m, 3H), 2.10 (m, 1H), 2.22 (m, 1H), 2.95 (s, 3H), 3.10-3.20 (m, 1H), 3.30-3.50 (m, 4H), 3.80 (m, 1H), 4.05 (m, 1H), 5.06 (s, 1H), 7.08-7.16 (m, 4H), 7.28-7.35 (m, 4H), 8.98 (d, 1H, NH).

Example 30

1-(3-Phenoxypropyl)-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane bromide

The title compound was synthesised according to methods a and d. The yield of the final step was 0.44 g, 100%.

m.p.: 242°C.

MS [M-Br]⁺: 469.

¹H-NMR (DMSO-d₆): δ 1.80-2.0 (m, 3H), 2.05-2.30 (m, 4H), 3.25-3.70 (m, 7H), 3.82 (m, 1H), 4.05 (t, 2H), 4.10 (m, 1H), 5.15 (s, 1H), 6.90-7.0 (m, 3H), 7.05-7.20 (m, 4H), 7.25-7.42 (m, 6H), 9.16 (d, 1H, NH).

Example 31

N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-9H-xanthene-9-carboxamide

The title compound was synthesised according to method a. The yield was 1.03 g, 43.4 %.

m.p.: 239.7-241.4°C.

MS [M+1]⁺: 335.

¹H-NMR (DMSO-d₆): δ 1.25-1.60 (m, 3H), 1.66 (m, 1H), 1.82 (m, 1H), 2.42-2.50 (m, 1H), 2.55-2.75 (m, 3H), 2.85 (m, 1H), 3.05 (m, 1H), 3.60 (m, 1H), 5.0 (s, 1H), 7.05-7.15 (m, 4H), 7.25-7.35 (m, 4H), 8.52 (d, 1H, NH).

5

Example 32**(3R)-1-(3-Phenoxypropyl)-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane bromide**

The title compound was synthesised according to methods a and d. The yield of the final
10 step was 0.48 g, 98%.

m.p.: 278.5-279.2°C.

MS [M-Br]⁺: 469.

¹H-NMR (DMSO-d₆): δ 1.80-2.0 (m, 3H), 2.05-2.30 (m, 4H), 3.15-3.25 (m, 1H), 3.30-3.65
(m, 6H), 3.82 (m, 1H), 4.05 (t, 2H), 4.10 (m, 1H), 5.05 (s, 1H), 6.90-7.0 (m, 3H), 7.05-7.20
15 (m, 4H), 7.25-7.42 (m, 6H), 9.01 (d, 1H, NH).

Example 33**N-[(3S)-1-Azabicyclo[2.2.2]oct-3-yl]-9H-xanthene-9-carboxamide**

The title compound was synthesised according to method a. The yield was 1.1 g, 60%.

20 m.p.: 244.2-244.9°C.

MS [M+1]⁺: 335.

¹H-NMR (DMSO-d₆): δ 1.25-1.60 (m, 3H), 1.66 (m, 1H), 1.82 (m, 1H), 2.42-2.50 (m, 1H),
2.55-2.75 (m, 3H), 2.83 (m, 1H), 3.05 (m, 1H), 3.60 (m, 1H), 5.0 (s, 1H), 7.05-7.15 (m,
4H), 7.25-7.35 (m, 4H), 8.53 (d, 1H, NH).

25

Example 34**(3S)-1-Allyl-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane trifluoroacetate**

The title compound was synthesised according to methods a and e. The yield of the final
30 step was 0.0091 g, 22.1%.

MS [M-CF₃COO]⁺: 375.

Example 35

**(3S)-1-Heptyl-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane
35 trifluoroacetate**

The title compound was synthesised according to methods a and e. The yield of the final step was 0.0097 g, 21.0 %.

MS [M-CF₃COO]⁺: 433.

5 **Example 36**

(3S)-1-Cyclohexylmethyl-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was synthesised according to methods a and e. The yield of the final step was 0.0124 g, 27.0 %.

10 MS [M-CF₃COO]⁺: 431.

Example 37

(3S)-1-(3-Cyclohexylpropyl)-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane trifluoroacetate

15 The title compound was synthesised according to methods a and e. The yield of the final step was 0.0045 g, 9.3%.

MS [M-CF₃COO]⁺: 459.

Example 38

20 **(3S)-1-(3-Phenoxypropyl)-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane bromide**

The title compound was synthesised according to methods a and d. The yield of the final step was 0.28 g, 85%.

m.p.: 279.0-280.4°C.

25 MS [M-Br]⁺: 469.

¹H-NMR (DMSO-d₆): δ 1.80-2.0 (m, 3H), 2.05-2.30 (m, 4H), 3.20-3.30 (m, 1H), 3.30-3.65 (m, 6H), 3.82 (m, 1H), 4.05 (t, 2H), 4.11 (m, 1H), 5.10 (s, 1H), 6.90-7.0 (m, 3H), 7.05-7.20 (m, 4H), 7.25-7.42 (m, 6H), 9.10 (d, 1H, NH).

30 **Example 39**

(3S)-1-[3-(5,6,7,8-Tetrahydronaphthalen-2-yloxy)propyl]-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane trifluoroacetate

The title compound was synthesised according to methods a and e. The yield of the final step was 0.0114 g, 21.2%.

35 MS [M-CF₃COO]⁺: 523.

Example 40

(3S)-1-[5-(2,6-Dimethylphenoxy)pentyl]-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane trifluoroacetate

- 5 The title compound was synthesised according to methods a and e. The yield of the final step was 0.0101 g, 18.7%.

MS [M-CF₃COO]⁺: 525.

Example 41

- 10 **N-(1-Azabicyclo[2.2.2]oct-3-yl)-9-hydroxy-9H-fluorene-9-carboxamide**

The title compound was synthesised according to method b. The yield was 1.06 g, 17.6%.

m.p.: 230°C.

MS [M+1]⁺: 335.

- 15 ¹H-NMR (CDCl₃): δ 1.0-1.30 (m, 2H), 1.52 (m, 2H), 1.79 (m, 1H), 2.04 (m, 1H), 2.36 (m, 1H), 2.57 (m, 3H), 3.05 (m, 1H), 3.77 (m, 1H), 5.45 (bs, 1H), 5.71 (d, 1H, NH), 7.20-7.50 (m, 6H), 7.55-7.70 (m, 2H).

Example 42

- 20 **3-[[[9-Hydroxy-9H-fluoren-9-yl)carbonyl]amino]-1-methyl-1-azoniabicyclo[2.2.2]octane bromide**

The title compound was synthesised according to methods b and d. The yield of the final step was 0.26 g, 83.8%.

m.p.: 197.5-203.6°C.

- 25 MS [M-Br]⁺: 349.

¹H-NMR (DMSO-d₆): δ 1.80-2.0 (m, 3H), 2.10-2.25 (m, 2H), 2.96 (s, 3H), 3.35-3.70 (m, 5H), 3.78 (m, 1H), 4.15 (m, 1H), 6.82 (s, 1H, OH), 7.30 (m, 2H), 7.43 (m, 4H), 7.79 (m, 2H), 8.81 (d, 1H, NH).

- 30 **Example 43**

3-[[[9-Hydroxy-9H-fluoren-9-yl)carbonyl]amino]-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane bromide

The title compound was synthesised according to methods b and d. The yield of the final step was 0.35 g, 87.5%.

- 35 m.p.: 264°C.

MS [M-Br]⁺: 469.

¹H-NMR (DMSO-d₆): δ 1.80-2.0 (m, 3H), 2.10-2.30 (m, 4H), 3.30-3.55 (m, 5H), 3.67 (m, 2H), 3.79 (m, 1H), 4.08 (t, 2H), 4.20 (m, 1H), 6.82 (s, 1H, OH), 6.97 (m, 3H), 7.32 (m, 4H), 7.43 (m, 4H), 7.79 (m, 2H), 8.88 (d, 1H, NH).

5

Example 44

N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-9-hydroxy-9H-fluorene-9-carboxamide

The title compound was synthesized according to method b. The yield was 0.304g, 4%.

MS [M+1]⁺: 335

10 ¹H-RMN (CDCl₃): δ 7.67 - 7.62 (m, 2H), 7.42 - 7.36 (m, 4H), 7.32 - 7.25 (m, 2H), 6.08 - 5.93 (m, NH, 1H), 3.74 (m, 1H), 3.00 - 2.87 (m, 1H), 2.52 - 2.27 (m, 4H), 2.02 - 1.92 (m, 1H), 1.80 - 1.76 (m, 1H), 1.51 - 1.44 (m, 2H), 1.26 - 1.17 (m, 2H).

Example 45

15 N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-fur-2-yl-2-hydroxypent-3-ynamide

The title compound was synthesized as a mixture of diastereomers according to method c. The yield was 1.38 g, 39%

MS [M+1]⁺: 289

20 ¹H-RMN (CDCl₃): (mixture of diastereomers) δ 7.36 - 7.35 (m, 1H), 6.87 - 6.76 (m, NH, 1H), 6.54 - 6.52 (m, 1H), 6.34 - 6.32 (m, 1H), 3.92 (m, 1H), 3.29 - 3.17 (m, 1H), 2.78 - 2.64 (m, 4H), 2.47 - 2.35 (m, 1H), 2.00 - 1.90 (m, 1H), 1.92 (s, 3H), 1.70 - 1.39 (m, 4H).

Example 46

25 N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-fur-2-yl-2-hydroxy-4-(4-methoxyphenyl)butanamide

The title compound was prepared as a mixture of diastereomers according to method c. The yield was 2.4 g, 44%

MS [M+1]⁺: 385

30 ¹H-RMN (CDCl₃): (mixture of diastereomers) δ 7.39 - 7.38 (m, 1H), 7.09 (d, 2H), 6.82 (d, 2H), 6.70 - 6.67 (m, NH, 1H), 6.41 - 6.35 (m, 2H), 3.92 (m, 1H), 3.78 (s, 3H), 3.33 - 3.24 (m, 1H), 2.82 - 2.19 (m, 9H), 1.96 - 1.87 (m, 1H), 1.67 - 1.40 (m, 4H)

Example 47

N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-N-methyl-9H-xanthene-9-carboxamide

35 The title product was synthesized according to method a. The yield was 0.9 g, 50%.

MS [M+1]⁺: 349

¹H-RMN (400 MHz, 60°C, CDCl₃): δ 7.26 - 7.19 (m, 4H), 7.10 - 7.01 (m, 4H), 5.51 (s, 1H), 4.18 (broad multiplet, 1H), 2.92 (s, 3H), 2.85-2.40 (m, 6H), 1.85-1.20 (m, 5H)

5 **Example 48**

N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-(5-bromothiophen-2-yl)-2-(4-fluoro-3-methylphenyl)-2-hydroxyacetamide

The title compound was synthesized as a mixture of diastereomers according to method c. The yield was 2.1 g, 42%.

10 MS [M+1]⁺: 453, 455

¹H-RMN (CDCl₃): (mixture of diastereomers) δ 7.35 - 7.20 (m, 2H), 7.08 - 6.89 (m, 3H), 6.80 (dd, 1H), 3.93 - 3.84 (m, 1H), 3.18 - 3.07 (m, 1H), 2.69 - 2.58 (m, 4H), 2.33 - 2.22 (m, 4H), 1.92 - 1.87 (m, 1H), 1.63 - 1.42 (m, 4H)

15 **Example 49**

(2S)-N-[(3R)-1-azabicyclo[2.2.2]oct-3-yl]-2-cyclopentyl-2-hydroxy-2-thien-2-ylacetamide

The title compound was synthesized according to method b. The yield was 0.650 g, 44%.

MS [M+1]⁺: 335

20 ¹H-RMN (CDCl₃): δ 7.24 (m, 1H), 7.08 (m, 1H), 6.97 (m, 1H), 6.64 (d, NH, 1H), 3.91 - 3.81 (m, 1H), 3.33 - 3.23 (m, 1H), 2.88 - 2.69 (m, 5H), 2.42 - 2.33 (m, 1H), 1.87 (m, 1H), 1.69 - 1.36 (m, 12H)

Example 50

25 **(3R)-1-[3-(Benzothiazol-2-yloxy)propyl]-3-(2-fur-2-yl-2-hydroxypent-3-ynoylamino)-1-azonia-bicyclo[2.2.2]octane chloride**

The title compound was synthesized as a mixture of diastereomers according to methods c and d.

30 The reaction time for the final step (conditions: THF, reflux temperature) was 7 days. The obtained product was purified by several washings with ether. The yield was 0.337 g (75%).

MS [M-Cl]⁺: 480

HPLC: mixture of diastereomers 49:51

¹H-NMR (DMSO-d₆): (mixture of diastereomers) δ 8.53 - 8.49 (m, NH, 1H), 7.69 (m, 1H), 7.58 (m, 1H), 7.53 - 7.39 (m, 2H), 7.26 - 7.19 (m, 2H), 6.43 - 6.38 (m, 2H), 4.20 - 4.14 (m, 1H), 4.05 - 3.99 (m, 2H), 3.76 - 3.65 (m, 1H), 3.60-3.20 (m, 7H), 2.20-1.65 (m, 7H), 1.87 (s, 3H)

5

Example 51

(3R)-3-(2-Fur-2-yl-2-hydroxypent-3-ynoylamino)-1-[3-(naphthalen-1-yloxy)propyl]-1-azoniabicyclo[2.2.2]octane chloride

The title compound was synthesized as a mixture of diastereomers according to methods c and d.

The reaction time for the final step (conditions, THF, reflux temperature) was 7 days. The obtained product was purified by several washings with ether. The yield was 0.252 g (57%).

MS [M-Cl]⁺: 473

15 HPLC: mixture of diastereomers 48:52

¹H-NMR (DMSO-d₆): (mixture of diastereomers) δ 8.62 - 8.58 (m, NH, 1H), 8.23 (m, 1H), 7.90 - 7.86 (m, 1H), 7.61 - 7.40 (m, 5H), 7.25 (m, 1H), 6.99 (m, 1H), 6.46 - 6.39 (m, 2H), 4.30 - 4.20 (m, 3H), 3.95 - 3.30 (m, 8H), 2.42 - 2.25 (m, 2H), 2.20 - 1.70 (m, 5H), 1.89 (s, 3H).

20

Example 52

(3R)-1-[3-(Benzo[1,3]dioxol-5-yloxy)propyl]-3-(2-fur-2-yl-2-hydroxypent-3-ynoylamino)-1-azonia-bicyclo[2.2.2]octane bromide

The title compound was synthesised as a mixture of diastereomers according to methods c and d.

The reaction time for the final step (conditions, THF, reflux temperature) was 2 days. The obtained product was purified by several washings with ether. The yield was 0.320 g (78%).

MS [M-Br]⁺: 467

30 HPLC: mixture of diastereomers 52:48

¹H-NMR (DMSO-d₆): (mixture of diastereomers) δ 8.54 - 8.49 (m, NH, 1H), 7.60 (s, 1H), 7.21 (d, 1H), 6.82 (d, 1H), 6.65 (m, 1H), 6.45 - 6.36 (m, 3H), 5.96 (s, 2H), 4.26 - 4.15 (m, 1H), 4.00 - 3.95 (m, 2H), 3.82 - 3.70 (m, 1H), 3.61 - 3.28 (m, 7H), 2.15 - 1.74 (m, 7H), 1.89 (s, 3H).

35

Example 53**(3R)-3-[[[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetyl]amino]-1-(4,4,4-trifluorobutyl)-1-azoniabicyclo[2.2.2]octane bromide**

The title compound was synthesised according to method d from Example 49.

- 5 The reaction time (conditions, THF, reflux temperature) was 3 days. The obtained product was purified by several washings with ether. The yield was 0.183 g (77%).

MS [M-Br]⁺: 445

- ¹H-NMR (DMSO-d₆): δ 8.28 (d, NH, 1H), 7.38 (dd, 1H), 7.09 (dd, 1H), 6.94 (dd, 1H), 6.11 (s, OH, 1H), 4.21 - 4.12 (m, 1H), 3.72 (m, 1H), 3.59 - 3.30 (m, 5H), 3.25 - 3.17 (m, 2H), 2.80 (m, 1H), 2.41 - 2.25 (m, 2H), 2.05 - 1.80 (m, 6H), 1.80-1.62 (m, 1H), 1.62-1.30 (m, 8H)
- 10

Example 54**(3R)-3-[[[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetyl]amino]-1-(2-hydroxyethyl)-1-azoniabicyclo[2.2.2]octane bromide**

The title compound was synthesised according to method d from Example 49.

The reaction time (conditions: THF, reflux temperature) was 5 days. The obtained product was purified by several washings with ether. The yield was 0.174 g (84%).

MS [M-Br]⁺: 379

- 20 ¹H-NMR (DMSO-d₆): δ 8.24 (d, NH, 1H), 7.37 (dd, 1H), 7.08 (dd, 1H), 6.94 (dd, 1H), 6.10 (s, OH, 1H), 5.27 (t, OH, 1H), 5.20-5.10 (m, 1H), 4.15 (m, 1H), 3.85 - 3.75 (m, 2H), 3.75-3.20 (m, 5H), 2.83 - 2.77 (m, 1H), 2.00 - 1.85 (m, 4H), 1.77 - 1.66 (m, 1H), 1.53 - 1.41 (m, 8H)

25 Example 55**(3R)-1-(4-Acetoxybutyl)-3-[2-(5-bromothien-2-yl)-2-(4-fluoro-3-methylphenyl)-2-hydroxyacetyl]amino]-1-azoniabicyclo[2.2.2]octane bromide**

The title compound was synthesised as a mixture of diastereomers according to methods c and d. The reaction time for the final step (conditions: THF, reflux temperature) was 41 hours. The obtained product was purified by several washings with ether. The yield was 0.190 g (88%).

30

MS [M-Br]⁺: 569, 567

¹H-NMR (DMSO-d₆): (mixture of diastereomers) δ 8.72 (d, NH, 1H), 7.45 (d, 1H), 7.36 - 7.18 (m, 2H), 7.13 - 7.06 (m, 2H), 6.91 (t, 1H), 4.25 - 4.18 (m, 1H), 4.06 - 4.00 (m, 2H),

3.72 - 3.26 (m, 6H), 3.19 - 3.11 (m, 2H), 2.20 (s, 3H), 2.21 - 1.84 (m, 1H), 2.02 (s, 3H),
1.95 - 1.54 (m, 8H)

Example 56

5 (3R)-3-[2-(5-Bromothiophen-2-yl)-2-(4-fluoro-3-methylphenyl)-2-hydroxyacetyl-amino]-1-(4-ethoxycarbonylbutyl)-1-azoniabicyclo[2.2.2]octane bromide

The title compound was synthesised as a mixture of diastereomers according to methods c and d. The reaction time for the final step (conditions: THF, reflux temperature) was 46 hours. The obtained product was purified by several washings with ether. The yield was

10 0.201 g (92%).

MS [M-Br]⁺: 583, 581

¹H-NMR (DMSO-d₆): (mixture of diastereomers) δ 8.72 (d, NH, 1H), 7.45 (d, 1H), 7.36 - 7.18 (m, 2H), 7.13 - 7.06 (m, 2H); 6.91 (t, 1H), 4.26 - 4.16 (m, 1H), 4.06 (q, 2H), 3.65 - 3.30 (m, 6H), 3.18 - 3.09 (m, 2H), 2.40 - 2.32 (m, 2H), 2.20 (s, 3H), 2.08 - 1.48 (m, 9H),

15 1.18 (t, 3H)

Example 57

(3R)-1-(3-Cyanopropyl)-3-[2-fur-2-yl-2-hydroxy-4-(4-methoxyphenyl)butyrylamino]-1-azoniabicyclo[2.2.2]octane bromide

20 The title compound was synthesised as a mixture of diastereomers according to methods c and d. The reaction time for the final step (conditions: THF, reflux temperature) was 3 days. The obtained product was purified by several washings with ether. The yield was 0.113 g (65%).

MS [M-Br]⁺: 452

25 ¹H-NMR (DMSO-d₆): (mixture of diastereomers) δ 8.49 - 8.40 (m, NH, 1H), 7.58 (s, 1H), 7.09 (d, 2H), 6.84 (d, 2H), 6.41 - 6.35 (m, 2H), 6.33 (d, 1H), 4.27 - 4.19 (m, 1H), 3.80-3.65 (m, 1H), 3.71 (s, 3H), 3.63 - 3.36 (m, 5H), 3.23 - 3.17 (m, 2H), 2.68 - 2.60 (m, 3H), 2.40 - 1.73 (m, 10H)

30 Example 58

(3R)-1-(2-[1,3]Dioxolan-2-yl-ethyl)-3-[2-fur-2-yl-2-hydroxy-4-(4-methoxyphenyl)butyrylamino]-1-azoniabicyclo[2.2.2]octane bromide

The title compound was synthesised as a mixture of diastereomers according to methods c and d. The reaction time for the final step (conditions: THF, reflux temperature) was 3

days. The obtained product was purified by several washings with ether. The yield was 0.128 g (70%).

MS [M-Br]⁺: 485

¹H-NMR (DMSO-d₆): (mixture of diastereomers) δ 8.47 - 8.37 (m, NH, 1H), 7.58 (s, 1H), 7.09 (d, 2H), 6.84 (d, 2H), 6.40 - 6.35 (m, 2H), 6.32 (d, 1H), 4.94 - 4.90 (m, 1H), 4.21 (m, 1H), 3.96 - 3.77 (m, 4H), 3.71 (s, 3H), 3.59 - 3.21 (m, 7H), 2.72 - 2.61 (m, 1H), 2.41 - 1.74 (m, 10H).

Example 59

10 (3R)-3-[[[9-Hydroxy-9H-fluoren-9-yl)carbonyl]amino]-1-[3-(methylphenylamino)propyl]-1-azoniabicyclo[2.2.2]octane chloride

The title compound was synthesised according to methods b and d. The reaction time for the final step (conditions: THF:CHCl₃ 1:1, reflux temperature) was 20 days. The obtained product was purified by washing with hot CHCl₃ and ether. The yield was 0.052 g (33%).

15 MS [M-Cl]⁺: 482

¹H-NMR (DMSO-d₆): δ 8.86 (d, NH, 1H), 7.78 (d, 2H), 7.45 - 7.16 (m, 8H), 6.81 - 6.62 (m, 4H), 4.23 - 4.14 (m, 1H), 3.76 - 3.53 (m, 2H), 3.43 - 3.17 (m, 8H), 2.91 (s, 3H), 2.23 - 1.77 (m, 7H).

20 Example 60

(3R)-3-[Methyl-(9H-xanthen-9-ylcarbonyl)amino]-1-(3-pyrrol-1-ylpropyl)-1-azoniabicyclo[2.2.2]octane bromide

The title compound was synthesised according to methods a and d. The reaction time for the final step (conditions: THF, reflux temperature) was 2 days. The obtained product was purified by several washings with THF and ether. The yield was 0.257 g (83%).

25 MS [M-Br]⁺: 456

¹H-NMR (DMSO-d₆): δ 7.34 - 7.06 (m, 8H), 6.78 (m, 2H), 6.01 (m, 2H), 5.75 (s, 1H), 4.47 (m, 1H), 3.94 - 3.88 (m, 2H), 3.72 - 3.25 (m, 6H), 3.39 (s, 3H), 3.20-3.05 (m, 2H), 2.22 - 1.88 (m, 7H).

30

Example 61

(3R)-1-[3-(Biphenyl-4-yloxy)propyl]-3-[methyl-(9H-xanthene-9-carbonyl)amino]-1-azoniabicyclo[2.2.2]octane chloride

The title compound was synthesised according to methods a and d. The reaction time for the final step (conditions: THF, reflux temperature) was 11 days. The obtained product was purified by several washings with THF and ether. The yield was 0.195 g (57%).

MS [M-Cl]⁺: 559

- 5 ¹H-NMR (DMSO-d₆): δ 7.62 (d, 4H), 7.47 - 7.02 (m, 13H), 5.77 (s, 1H), 4.50 (m, 1H), 4.13 - 4.06 (m, 2H), 3.82 - 3.31 (m, 8H), 3.42 (s, 3H), 2.25 - 1.91 (m, 7H).

- 10 The following examples illustrate pharmaceutical compositions according to the present invention and procedures for their preparation.

Example 62

Preparation of a pharmaceutical composition: tablets

Formulation:

- | | | |
|----|-----------------------------------|----------|
| 15 | Compound of the present invention | 5.0 mg |
| | Lactose | 113.6 mg |
| | Microcrystalline cellulose | 28.4 mg |
| | Light silicic anhydride | 1.5 mg |
| | Magnesium stearate | 1.5 mg |

20

Using a mixer machine, 15 g of the compound of the present invention was mixed with 340.8 g of lactose and 85.2 g of microcrystalline cellulose. The mixture was subjected to compression moulding using a roller compactor to give a flake-like compressed material. The flake-like compressed material was pulverized using a hammer mill, and the

25 pulverized material was screened through a 20 mesh screen. A 4.5 g portion of light silicic anhydride and 4.5 g of magnesium stearate were added to the screened material and mixed. The mixer product was subjected to a tablets making machine equipped with a die/punch system of 7.5 mm in diameter, thereby obtaining 3,000 tablets each having 150 mg in weight.

30

Example 63

Preparation of a pharmaceutical composition: coated tablets

Formulation:

- | | | |
|----|-----------------------------------|---------|
| | Compound of the present invention | 5.0 mg |
| 35 | Lactose | 95.2 mg |

	Corn starch	40.8 mg
	Polyvinylpyrrolidone K25	7.5 mg
	Magnesium stearate	1.5 mg
	Hydroxypropylcellulose	2.3 mg
5	Polyethylene glycol 6000	0.4 mg
	Titanium dioxide	1.1 mg
	Purified talc	0.7 mg

Using a fluidized bed granulating machine, 15 g of the compound of the present invention
 10 was mixed with 285.6 g of lactose and 122.4 g of corn starch. Separately, 22.5 g of polyvinylpyrrolidone was dissolved in 127.5 g of water to prepare a binding solution. Using a fluidized bed granulating machine, the binding solution was sprayed on the above mixture to give granulates. A 4.5 g portion of magnesium stearate was added to the obtained granulates and mixed. The obtained mixture was subjected to a tablet making
 15 machine equipped with a die/punch biconcave system of 6.5 mm in diameter, thereby obtaining 3,000 tablets, each having 150 mg in weight.

Separately, a coating solution was prepared by suspending 6.9 g of hydroxypropylmethylcellulose 2910, 1.2 g of polyethylene glycol 6000, 3.3 g of titanium dioxide and 2.1 g of purified talc in 72.6 g of water. Using a High Coated, the 3,000 tablets
 20 prepared above were coated with the coating solution to give film-coated tablets, each having 154.5 mg in weight.

Example 64

Preparation of a pharmaceutical composition: liquid inhalant

25 Formulation:

Compound of the present invention	400 µg
Physiological saline	1 ml

A 40 mg portion of the compound of the present invention was dissolved in 90 ml of
 30 physiological saline, and the solution was adjusted to a total volume of 100 ml with the same saline solution, dispensed in 1 ml portions into 1 ml capacity ampoule and then sterilized at 115 °C for 30 minutes to give a liquid inhalant.

Example 65

35 Preparation of a pharmaceutical composition: powder inhalant

Formulation:

Compound of the present invention	200 µg
Lactose	4,000 µg

- 5 A 20 g portion of the compound of the present invention was uniformly mixed with 400 g of lactose, and a 200 mg portion of the mixture was packed in a powder inhaler for exclusive use to produce a powder inhalant.

Example 66**10 Preparation of a pharmaceutical composition: inhalation aerosol.****Formulation:**


Compound of the present invention	200 µg
Dehydrated (Absolute) ethyl alcohol USP	8,400 µg
1,1,1,2-Tetrafluoroethane (HFC-134A)	46,810 µg

15

The active ingredient concentrate is prepared by dissolving 0.0480 g of the compound of the present invention in 2.0160 g of ethyl alcohol. The concentrate is added to an appropriate filling apparatus. The active ingredient concentrate is dispensed into aerosol container, the headspace of the container is purged with Nitrogen or HFC-134A vapour
20 (purging ingredients should not contain more than 1 ppm oxygen) and is sealed with valve. 11.2344 g of HFC-134A propellant is then pressure filled into the sealed container.

CLAIMS

1. A compound which is a quinuclidine amide derivative of formula (I):



(I)


wherein

R₁ represents a hydrogen atom or an straight or branched, optionally substituted lower


alkyl group;

R₂ represents a group of formula i) or ii)

i)



ii)



wherein

R₃ represents a group selected from phenyl, 2-furyl, 3-furyl, 2-thienyl or 3-thienyl;

R₄ represents a group selected from optionally substituted lower alkyl, optionally

substituted lower alkenyl, optionally substituted lower alkynyl, cycloalkyl, cycloalkylmethyl, phenyl, benzyl, phenethyl, 2-furyl, 3-furyl, 2-thienyl or 3-thienyl;

and R₅ represents a hydrogen atom or a hydroxy, methyl, or -CH₂OH group;

the benzene rings in formula ii) and the cyclic groups represented by R₃ and R₄ being each independently optionally substituted by one, two or three substituents selected from

halogen, straight or branched, optionally substituted lower alkyl, hydroxy, straight or branched, optionally substituted lower alkoxy, nitro, cyano, $-\text{CO}_2\text{R}'$ or $-\text{NR}'\text{R}''$, wherein R' and R'' each independently represents a hydrogen atom or a straight or branched, optionally substituted lower alkyl group or R' and R'' together with the atom to which they are attached form a cyclic group;

Q represents a single bond or a $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, $-\text{O}-$, $-\text{OCH}_2-$, $-\text{S}-$, $-\text{SCH}_2-$ or $-\text{CH}=\text{CH}-$ group;

p is 1 or 2 and the amide group is at positions 2, 3 or 4 of the azabicyclic ring;

or pharmaceutically acceptable salts thereof, including quaternary ammonium salts;


and all individual stereoisomers and mixtures thereof;

with the proviso that when p is 2, the amide moiety is in position 3 of the quinuclidine ring, R_1 is hydrogen and R_3 and R_4 are both unsubstituted phenyl, then

-when said compound is not a pharmaceutically acceptable salt or is a HCl salt, then R_5 cannot be one of hydrogen or hydroxy; and

- when said compound is a quaternary ammonium salt having a methyl group attached to the nitrogen atom of the quinuclidine ring, then R_5 cannot be hydroxy.

2. A compound according to claim 1 which is a quaternary ammonium salt of formula (II)



(II)

wherein R_1 , R_2 and p are as defined in claim 1;

m is an integer from 0 to 8;

n is an integer from 0 to 4;

A represents a group selected from $-\text{CH}_2-$, $-\text{CH}=\text{CR}'-$, $-\text{CR}'=\text{CH}-$, $-\text{CR}'\text{R}''-$, $-\text{C}(\text{O})-$, $-\text{O}-$, $-\text{S}-$, $-\text{S}(\text{O})-$, $-\text{S}(\text{O})_2-$ and $-\text{NR}'-$, wherein R' and R'' are as defined in claim 1;

- 5 R_6 represents a hydrogen atom, or a group selected from straight or branched, optionally substituted lower alkyl, hydroxy, straight or branched, optionally substituted lower alkoxy, cyano, nitro, $-\text{CH}=\text{CR}'\text{R}''$, $-\text{C}(\text{O})\text{OR}'$, $-\text{OC}(\text{O})\text{R}'$, $-\text{SC}(\text{O})\text{R}'$, $-\text{C}(\text{O})\text{NR}'\text{R}''$, $-\text{NR}'\text{C}(\text{O})\text{OR}''$, $-\text{NR}'\text{C}(\text{O})\text{NR}''$, cycloalkyl, phenyl, naphthanelyl, 5,6,7,8-tetrahydronaphthanelyl, benzo[1,3]dioxolyl, heteroaryl or heterocyclyl; R' and R'' being as defined in claim 1; and
- 10 wherein the cyclic groups represented by R_6 are optionally substituted by one, two or three substituents selected from halogen, hydroxy, straight or branched, optionally substituted lower alkyl, phenyl, $-\text{OR}'$, $-\text{SR}'$, $-\text{NR}'\text{R}''$, $-\text{NHCOR}'$, $-\text{CONR}'\text{R}''$, $-\text{CN}$, $-\text{NO}_2$ and $-\text{COOR}'$; R' and R'' being as defined in claim 1; and
- 15 X^- represents a pharmaceutically acceptable anion of a mono or polyvalent acid.

and all individual stereoisomers and mixtures thereof;

with the proviso that when p is 2, the amide moiety is in position 3 of the quinuclidine ring,

- 20 R_1 is hydrogen, R_3 and R_4 are both unsubstituted phenyl and R_5 is hydroxy, then in the compounds of formula (II) the sequence $\text{R}_6 - (\text{CH}_2)_n - \text{A} - (\text{CH}_2)_m -$ cannot be a methyl group.

3. A compound according to claim 1 or claim 2 wherein R_1 is hydrogen, methyl or ethyl.

25

4. A compound according to claim 3, wherein R_1 is hydrogen.

5. A compound according to any one of the preceding claims, wherein R_2 is a group of formula i), wherein R_3 is a group, which is optionally substituted with one or more halogen

30 atom(s), selected from phenyl, 2-thienyl, 3-thienyl or 2-furyl

6. A compound according to claim 5, wherein R_2 is a group of formula i), wherein R_3 represents a group phenyl, 2-thienyl or 2-furyl which are optionally substituted with one or more halogen atom(s).

35

7. A compound according to any one of claims 5 and 6, wherein R_2 is a group of formula i), wherein R_3 represents phenyl or 2-thienyl.

5 8. A compound according to any one of the preceding claims, wherein R_4 represents a linear group selected from ethyl, n-butyl, vinyl, allyl, 1-propenyl and 1-propynyl, or a group, which is optionally substituted with one or more halogen atom(s), methyl or methoxy group(s), selected from cyclopentyl, cyclohexyl, phenyl, benzyl, phenethyl, 2-thienyl and 3-furyl.

10

9. A compound according to claim 8, wherein R_4 represents a linear group selected from ethyl, n-butyl, vinyl, allyl and 1-propynyl, or a group, which is optionally substituted with one or more halogen atom(s), methyl or methoxy group(s), selected from cyclopentyl, phenyl, benzyl, phenethyl and 2-thienyl.

15

10. A compound according to any one of claims 8 and 9, wherein R_4 represents a group selected from ethyl, n-butyl, vinyl, allyl, cyclopentyl, phenyl, benzyl or 2-thienyl

20

11. A compound according to any one of claims 1 to 4, wherein R_2 is a group of formula ii), wherein Q represents a single bond or an oxygen atom.

12. A compound according to any one of the preceding claims wherein R_5 is hydrogen or hydroxy.

25 13. A compound according to any one of the preceding claims wherein p is 2 and the amide group is at positions 3 or 4 of the azabicyclic ring.

14. A compound according to claim 13 wherein the amide group is at position 3 of the azabicyclic ring.

30

15. A compound of formula (II) according to any one of claims 2 to 14, wherein m is an integer from 0 to 6 and n is an integer from 0 to 4; A represents a group selected from $-CH_2-$, $-CH=CH-$, $-O-$, $-C(O)-$, $-NR'-$, and $-S-$; and R_6 is a hydrogen atom, a cyano group, a nitro group, a $-C(O)OR'$ group, a $-OC(O)R'$ group, a $-SC(O)R'$ group, a $-CH=CH_2$ group, 35 a $-CH=CR'R''$ group, a $C(O)NR'R''$ group, a straight or branched C_1-C_4 alkyl group, which

is optionally substituted with one or more halogen atom(s), a straight C₁-C₄ alkoxy group, which is optionally substituted with one or more halogen atom(s) or hydroxy group(s), or a cyclic group, which is optionally substituted with one or more substituents selected from halogen atoms, groups of formula -C(O)NR'R'' and methyl, hydroxy, nitro and phenyl groups, the cyclic group being selected from cyclohexyl, phenyl, 5,6,7,8-tetrahydronaphthanyl, 2-thienyl, 1-pyrrolidinyl, 1-pyrrolyl, benzo[1,3]dioxolyl, 2-benzothiazolyl, naphthalenyl and dioxolyl.

16. A compound according to claim 15, wherein m is an integer from 0 to 5 and n is an integer from 0 to 2; A represents a group selected from -CH₂-, -CH=CH-, -O-, -C(O)-, -NR'-, and -S-; and R₆ is a hydrogen atom, a cyano group, a -C(O)OR' group, a -OC(O)R' group, a -SC(O)R' group, a -CH=CH₂ group, a -C(O)NR'R'' group, a straight or branched C₁-C₄ alkyl group, a trifluoromethyl, or a cyclic group selected from cyclohexyl, 5,6,7,8-tetrahydronaphthanyl, 2-thienyl, 1-pyrrolyl, benzo[1,3]dioxolyl, 2-benzothiazolyl, naphthalenyl, dioxolyl and phenyl, which is optionally substituted with one or more substituents selected from halogen atoms, groups of formula -C(O)NR'R'', methyl, hydroxy and phenyl groups.

17. A compound according to any one of claims 15 and 16, wherein m is an integer from 0 to 5 and n is an integer from 0 to 2; A represents a group selected from -CH₂-, -CH=CH-, -O-; and R₆ is selected from hydrogen, straight C₁-C₄ alkyl, -CH=CH₂, cyclohexyl, phenyl which is unsubstituted or substituted with one or two substituents selected from methyl groups and hydroxy groups, 5,6,7,8-tetrahydronaphthanyl and 2-thienyl.

18. A compound according to claim 17, wherein the sequence R₆ - (CH₂)_n - A - (CH₂)_m - is one of methyl, 3-phenoxypropyl, 3-(3-hydroxyphenoxy)propyl, allyl, heptyl, 3-phenylpropyl, 3-phenylallyl, 2-phenoxyethyl, 2-benzyloxyethyl, cyclohexylmethyl, 3-(5,6,7,8-tetrahydronaphthalen-2-yloxy)propyl, 5-(2,6-dimethylphenoxy)pentyl, 3-thien-2-ylpropyl or 3-cyclohexylpropyl and X' is bromide or trifluoroacetate.

19. A compound according to any one of the preceding claims, which is a single isomer.

20. A compound according to claim 1 or claim 2 which is one of:

N-(1-Azabicyclo[2.2.2]oct-3-yl)-2-hydroxy-2,2-dithien-2-ylacetamide

N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2,2-dithien-2-ylacetamide

- N-[(3S)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2,2-dithien-2-ylacetamide
 N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2,2-dithien-2-ylacetamide
 N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-cyclopentyl-2-hydroxy-2-thien-2-ylacetamide
 N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2-thien-2-ylpent-4-enamide
 5 (2*)-N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2-thien-2-ylbutanamide
 (diastereomer 1)
 (2*)-N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2-thien-2-ylbutanamide
 (diastereomer 2)
 (2*)-N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2-thien-2-ylbut-3-enamide
 10 (diastereomer 1)
 (2*)-N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-hydroxy-2-thien-2-ylbut-3-enamide
 (diastereomer 2)
 N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2,3-diphenylpropanamide
 N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-cyclopentyl-2-hydroxy-2-phenylacetamide
 15 N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-phenylhexanamide
 N-(1-Azabicyclo[2.2.2]oct-3-yl)-9H-xanthene-9-carboxamide
 N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-9H-xanthene-9-carboxamide
 N-[(3S)-1-Azabicyclo[2.2.2]oct-3-yl]-9H-xanthene-9-carboxamide
 N-(1-Azabicyclo[2.2.2]oct-3-yl)-9-hydroxy-9H-fluorene-9-carboxamide
 20 N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-N-methyl-9H-xanthene-9-carboxamide
 (2S)-N-[(3R)-1-azabicyclo[2.2.2]oct-3-yl]-2-cyclopentyl-2-hydroxy-2-thien-2-ylacetamide
 3-(2-Hydroxy-2,2-dithien-2-ylacetyl-amino)-1-methyl-1-azoniabicyclo[2.2.2]octane bromide
 3-(2-Hydroxy-2,2-dithien-2-ylacetyl-amino)-1-(3-phenoxypropyl)-1-
 azoniabicyclo[2.2.2]octane bromide
 25 (3R)-3-(2-Hydroxy-2,2-dithien-2-ylacetyl-amino)-1-methyl-1-azoniabicyclo[2.2.2]octane
 trifluoroacetate
 (3R)-1-Allyl-3-(2-hydroxy-2,2-dithien-2-ylacetyl-amino)-1-azoniabicyclo[2.2.2]octane
 trifluoroacetate
 (3R)-1-Heptyl-3-(2-hydroxy-2,2-dithien-2-ylacetyl-amino)-1-azoniabicyclo[2.2.2]octane
 30 trifluoroacetate
 (3R)-3-(2-Hydroxy-2,2-dithien-2-ylacetyl-amino)-1-(3-phenylpropyl)-1-
 azoniabicyclo[2.2.2]octane bromide
 (3R)-3-(2-Hydroxy-2,2-dithien-2-ylacetyl-amino)-1-((E)-3-phenylallyl)-1-
 azoniabicyclo[2.2.2]octane trifluoroacetate

- (3R)-3-(2-Hydroxy-2,2-dithien-2-ylacetyl amino)-1-(2-phenoxyethyl)-1-azoniabicyclo[2.2.2]octane bromide
- (3R)-3-(2-Hydroxy-2,2-dithien-2-ylacetyl amino)-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane bromide
- 5 (3R)-3-(2-Hydroxy-2,2-dithien-2-ylacetyl amino)-1-[3-(3-hydroxyphenoxy)propyl]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-1-(2-Benzoyloxyethyl)-3-(2-hydroxy-2,2-dithien-2-ylacetyl amino)-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3R)-3-(2-Hydroxy-2,2-dithien-2-ylacetyl amino)-1-(3-thien-2-ylpropyl)-1-azoniabicyclo[2.2.2]octane bromide
- 10 (3S)-3-(2-Hydroxy-2,2-dithien-2-ylacetyl amino)-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane bromide
- (3R)-3-(2,2-Dithien-2-ylacetyl amino)-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane bromide
- 15 1-Methyl-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane bromide
- 1-(3-Phenoxypropyl)-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane bromide
- (3R)-1-(3-Phenoxypropyl)-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane bromide
- 20 (3S)-1-Allyl-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3S)-1-Heptyl-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3S)-1-Cyclohexylmethyl-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 25 (3S)-1-(3-Cyclohexylpropyl)-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3S)-1-(3-Phenoxypropyl)-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane bromide
- 30 (3S)-1-[3-(5,6,7,8-Tetrahydronaphthalen-2-yloxy)propyl]-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- (3S)-1-[5-(2,6-Dimethylphenoxy)pentyl]-3-[(9H-xanthen-9-ylcarbonyl)amino]-1-azoniabicyclo[2.2.2]octane trifluoroacetate
- 35 3-[(9-Hydroxy-9H-fluoren-9-yl)carbonyl]amino-1-methyl-1-azoniabicyclo[2.2.2]octane bromide

- 3-[[[(9-Hydroxy-9H-fluoren-9-yl)carbonyl]amino]-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane bromide
 (3R)-1-[3-(2-Carbamoylphenoxy)propyl]-3-[[[(9-Hydroxy-9H-fluoren-9-yl)carbonyl]amino]-1-azoniabicyclo[2.2.2]octane formate
- 5 (3R)-1-[4-(4-Fluorophenyl)-4-oxobutyl]-3-[[[(9-Hydroxy-9H-fluoren-9-yl)carbonyl]amino]-1-azoniabicyclo[2.2.2]octane formate
 (3R)-3-[[[(9-Hydroxy-9H-fluoren-9-yl)carbonyl]amino]-1-[3-(methylphenylamino)propyl]-1-azoniabicyclo[2.2.2]octane chloride
 (3R)-3-[[[(9-Hydroxy-9H-fluoren-9-yl)carbonyl]amino]-1-(3-phenylsulfanylpropyl)-1-azoniabicyclo[2.2.2]octane formate
- 10 (3R)-3-[Methyl-(9H-xanthen-9-ylcarbonyl)amino]-1-(3-pyrrol-1-ylpropyl)-1-azoniabicyclo[2.2.2]octane bromide
 (3R)-1-[3-(Biphenyl-4-yloxy)propyl]-3-[methyl-(9H-xanthene-9-carbonyl)amino]-1-azoniabicyclo[2.2.2]octane chloride
- 15 (3R)-3-(2-Fur-2-yl-2-hydroxypent-3-ynoylamino)-1-[3-(naphthalen-1-yloxy)propyl]-1-azoniabicyclo[2.2.2]octane chloride
 (3R)-1-[3-(Benzo[1,3]dioxol-5-yloxy)propyl]-3-(2-fur-2-yl-2-hydroxypent-3-ynoylamino)-1-azonia-bicyclo[2.2.2]octane bromide
 (3R)-1-[3-(Benzothiazol-2-yloxy)propyl]-3-(2-fur-2-yl-2-hydroxypent-3-ynoylamino)-1-azonia-bicyclo[2.2.2]octane chloride
- 20 (3R)-3-[[[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetyl]amino]-1-(2-hydroxyethyl)-1-azoniabicyclo[2.2.2]octane bromide
 (3R)-3-[[[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetyl]amino]-1-(2-ethoxyethyl)-1-azoniabicyclo[2.2.2]octane formate
- 25 (3R)-3-[[[(2S)-2-Cyclopentyl-2-hydroxy-2-thien-2-ylacetyl]amino]-1-(4,4,4-trifluorobutyl)-1-azoniabicyclo[2.2.2]octane bromide
 (3R)-1-(4-Acetoxybutyl)-3-[2-(5-bromothien-2-yl)-2-(4-fluoro-3-methylphenyl)-2-hydroxyacetyl]amino]-1-azoniabicyclo[2.2.2]octane bromide
 (3R)-3-[2-(5-Bromothien-2-yl)-2-(4-fluoro-3-methylphenyl)-2-hydroxyacetyl]amino]-1-(4-ethoxycarbonylbutyl)-1-azoniabicyclo[2.2.2]octane bromide
- 30 (3R)-1-(3-Acetylsulfanylpropyl)-3-[2-(5-bromothien-2-yl)-2-(4-fluoro-3-methylphenyl)-2-hydroxyacetyl]amino]-1-azoniabicyclo[2.2.2]octane formate
 (3R)-1-(3-Cyanopropyl)-3-[2-fur-2-yl-2-hydroxy-4-(4-methoxyphenyl)butyrylamino]-1-azoniabicyclo[2.2.2]octane bromide


(3R)-1-(2-Carbamoylethyl)-3-[2-fur-2-yl-2-hydroxy-4-(4-methoxyphenyl)butyrylamino]-1-azoniabicyclo[2.2.2]octane formate

(3R)-1-(2-[1,3]Dioxolan-2-yl-ethyl)-3-[2-fur-2-yl-2-hydroxy-4-(4-methoxyphenyl)butyrylamino]-1-azoniabicyclo[2.2.2]octane bromide

5

21. A process for producing a compound of formula (I), as defined in claim 1 and wherein R_2 is a group of formula i) and R_5 is an hydroxy group, which process comprises reacting a compound of formula (V).

10



(V)

wherein R_1 and R_3 are as defined in claim 1 with the corresponding organometallic derivative R_4 -[Mg,Li], wherein R_4 is as defined in claim 1.

22. A compound of formula (V), which is one of

15

N-1-azabicyclo[2.2.2]oct-3-yl-2-oxo-2-thien-2-ylacetamide

N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-2-oxo-2-thien-2-ylacetamide

N-[(3S)-1-Azabicyclo[2.2.2]oct-3-yl]-2-oxo-2-thien-2-ylacetamide

20 23. A process for producing a compound of formula (II), as defined in claim 2, which process comprises reacting a compound of formula (I), as defined in claim 1 with an alkylating agent of formula $R_6-(CH_2)_n-A-(CH_2)_m-W$, wherein R_6 , n , A and m are as defined in claim 2 and W represents any suitable leaving group.

25 24. A pharmaceutical composition comprising a compound according to any one of claims 1 to 20 in admixture with a pharmaceutically acceptable carrier or diluent.

25. A compound according to any one of claims 1 to 20 for the treatment of a pathological condition or disease susceptible to amelioration by antagonism of M3 muscarinic receptors.
- 5 26. Use of a compound according to any one of claims 1 to 20 in the manufacture of a medicament for the treatment of a pathological condition or disease susceptible to amelioration by antagonism of M3 muscarinic receptors.
27. Use according to claim 26 wherein the pathological condition is a respiratory,
10 urological or gastrointestinal disease or disorder.
28. A method for treating a subject afflicted with a pathological condition or disease susceptible to amelioration by antagonism of M3 muscarinic receptors, which comprises administering to said subject an effective amount of a compound as defined in any one of
15 claims 1 to 20.
29. A method according to claim 28 wherein the pathological condition is a respiratory, urological or gastrointestinal disease or disorder.
- 20 30. A combination product comprising
(i) a compound according to any one of claims 1 to 20; and
(ii) another compound effective in the treatment of a respiratory, urological or gastrointestinal disease or disorder
for simultaneous, separate or sequential use.
- 25 31. A combination product according to claim 30 comprising
(i) a compound according to any one of claims 1 to 20; and
(ii) a β_2 agonist, steroid, antiallergic drug, phosphodiesterase IV inhibitor and/or leukotriene D4 (LTD4) antagonist for simultaneous, separate or sequential use in the
30 treatment of a respiratory disease.

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D453/02 C07D487/08 A61K31/439 A61K31/407 A61P1/00
A61P11/00 A61P13/00 //(C07D487/08,209:00,209:00)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	CA 2 155 320 A (FUJISAWA PHARMACEUTICAL CO) 19 August 1993 (1993-08-19) cited in the application page 1; claim 1 ---	1,2,24, 27
A	WO 01 04118 A (ALMIRALL PRODESFARMA SA) 18 January 2001 (2001-01-18) page 1, line 7 - line 18; claims 1,31,33 ---	1,2,24, 25,27
A	EP 0 930 298 A (BANYU PHARMA CO LTD) 21 July 1999 (1999-07-21) claims 1,15 ---	1,2,24, 25,27
P,A	WO 02 051841 A (ALMIRALL PRODESFARMA) 4 July 2002 (2002-07-04) claims 1,22,23,25 --- -/--	1,2,25, 27

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *S* document member of the same patent family

Date of the actual completion of the international search

1 October 2003

Date of mailing of the international search report

09/10/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Alfaro Faus, I

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 03/06708

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, A	WO 02 053564 A (ALMIRALL PRODESFARMA) 11 July 2002 (2002-07-11) claims 1,31,33 -----	1,2,25, 27

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 03/06708

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
CA 2155320	A	19-08-1993	CA 2155320 A1	19-08-1993
			WO 9316048 A1	19-08-1993
WO 0104118	A	18-01-2001	ES 2165768 A1	16-03-2002
			AT 235492 T	15-04-2003
			AU 6433000 A	30-01-2001
			BG 106301 A	30-08-2002
			BR 0012434 A	02-04-2002
			CA 2381165 A1	18-01-2001
			CN 1373760 T	09-10-2002
			CZ 20020121 A3	13-11-2002
			DE 60001840 D1	30-04-2003
			DK 1200431 T3	21-07-2003
			EE 200200017 A	15-04-2003
			WO 0104118 A2	18-01-2001
			EP 1200431 A2	02-05-2002
			HU 0202100 A2	28-10-2002
			JP 2003504368 T	04-02-2003
			NO 20020180 A	13-03-2002
			SK 432002 A3	01-04-2003
			TR 200200768 T2	22-07-2002
			US 2003055080 A1	20-03-2003
EP 0930298	A	21-07-1999	AT 229941 T	15-01-2003
			AU 716050 B2	17-02-2000
			AU 3635197 A	25-02-1998
			BG 103114 A	30-11-1999
			BR 9711108 A	17-08-1999
			DE 69718026 D1	30-01-2003
			DE 69718026 T2	10-07-2003
			EE 9900038 A	16-08-1999
			EP 0930298 A1	21-07-1999
			JP 3063164 B2	12-07-2000
			NO 990472 A	01-02-1999
			NZ 333842 A	25-05-2001
			SK 12299 A3	16-05-2000
			CN 1226888 A	25-08-1999
			CZ 9900331 A3	14-07-1999
			ES 2188961 T3	01-07-2003
			HR 970426 A1	31-08-1998
			HU 9902381 A2	29-11-1999
			WO 9805641 A1	12-02-1998
			JP 3282617 B2	20-05-2002
			JP 2000178231 A	27-06-2000
			JP 3282618 B2	20-05-2002
			JP 2000169449 A	20-06-2000
			KR 2000022214 A	25-04-2000
			PL 331431 A1	19-07-1999
			TR 9900204 T2	21-01-2000
			TR 200001482 T2	21-11-2000
			US 6040449 A	21-03-2000
			US 5948792 A	07-09-1999
			ZA 9706813 A	11-02-1998
WO 02051841	A	04-07-2002	WO 02051841 A1	04-07-2002
			EP 1345937 A1	24-09-2003
			NO 20032889 A	08-08-2003

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 03/06708

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 02053564	A	11-07-2002	CA 2433128 A1	11-07-2002
			WO 02053564 A2	11-07-2002
			NO 20033002 A	30-06-2003
<hr/>				